

OPERATORS RELATED TO SUBORDINATION FOR FREE MULTIPLICATIVE CONVOLUTIONS

ROMUALD LENCZEWSKI

ABSTRACT. It has been shown by Voiculescu and Biane that the analytic subordination property holds for free additive and multiplicative convolutions. In this paper, we present an operatorial approach to subordination for free multiplicative convolutions. This study is based on the concepts of ‘freeness with subordination’, or ‘s-free independence’, and ‘orthogonal independence’, introduced recently in the context of free additive convolutions. In particular, we introduce and study the associated multiplicative convolutions and construct related operators, called ‘subordination operators’ and ‘subordination branches’. Using orthogonal independence, we derive decompositions of subordination branches and related decompositions of s-free and free multiplicative convolutions. The operatorial methods lead to several new types of graph products, called ‘loop products’, associated with different notions of independence (monotone, boolean, orthogonal, s-free). We also prove that the enumeration of rooted ‘alternating double return walks’ on the loop products of graphs and on the free product of graphs gives the moments of the corresponding multiplicative convolutions.

1. INTRODUCTION

Multiplication of free random variables X_1 and X_2 with distributions μ_1 and μ_2 , respectively, leads to the multiplicative convolution $\mu_1 \boxtimes \mu_2$, introduced by Voiculescu [23] in the C^* -algebra framework, which gives the distribution of the product of X_1 and X_2 (the general case of measures with unbounded support was studied by Bercovici and Voiculescu [7]).

Let $\mathcal{M}_{\mathbb{R}_+}$ denote the set of probability measures on $\mathbb{R}_+ = [0, \infty)$. If $\mu \in \mathcal{M}_{\mathbb{R}_+}$, we can define

$$(1.1) \quad \psi_\mu(z) = \int_{\mathbb{R}_+} \frac{zt}{1-zt} d\mu(t), \quad z \in \mathbb{C} \setminus \mathbb{R}_+,$$

which, in the case when μ has finite moments of all orders, becomes the moment generating function $\psi_\mu(z) = \sum_{n=1}^{\infty} \mu(X^n) z^n$, where $\mu(X^n)$ are the moments of the unique functional $\mu : \mathbb{C}[X] \rightarrow \mathbb{C}$ defined by μ . In order to study $\mu_1 \boxtimes \mu_2$, Voiculescu introduced the S-transform of μ defined by $S_\mu(z) = (1+z)\psi_\mu^{-1}(z)/z$, where $\psi_\mu^{-1}(z)$ denotes the inverse of $\psi_\mu(z)$ with respect to composition. The key multiplicative formula for the S-transforms is given by $S_{\mu_1 \boxtimes \mu_2}(z) = S_{\mu_1}(z)S_{\mu_2}(z)$.

2000 Mathematics Subject Classification: Primary 46L54, 46L53; Secondary 05C50

Key words and phrases: free probability, free random variable, free multiplicative convolution, s-free multiplicative convolution, orthogonal multiplicative convolution, s-free independence, subordination, free product of graphs

This work is partially supported by MNiSW research grant No 1 P03A 013 30

In our approach, a central role is played by the transform related to $\psi_\mu(z)$, namely

$$(1.2) \quad \eta_\mu(z) = \frac{\psi_\mu(z)}{1 + \psi_\mu(z)}$$

where $z \in \mathbb{C} \setminus \mathbb{R}_+$. Using this transform, Biane [8] has proved the subordination property for free multiplicative convolutions of probability measures on $\mathbb{R}_+$ (and also for probability measures on the unit circle $\mathbb{T}$). Earlier, the subordination property for free additive convolutions [22] was discovered by Voiculescu [24] for compactly supported measures on $\mathbb{R}$, generalized by Biane [8] to arbitrary measures on $\mathbb{R}$ (see also [17] for a related approach).

For instance, subordination for free multiplicative convolutions of probability measures on $\mathbb{R}_+$ says that for given $\mu_1, \mu_2 \in \mathcal{M}_{\mathbb{R}_+} \setminus \{\delta_0\}$, there exist analytic self-maps η_1, η_2 of $\mathbb{C} \setminus \mathbb{R}_+$, such that

$$(1.3) \quad \eta_{\mu_1 \boxtimes \mu_2}(z) = \eta_{\mu_1}(\eta_1(z)) = \eta_{\mu_2}(\eta_2(z)).$$

One says that $\eta_{\mu_1 \boxtimes \mu_2}$ is *subordinate* to both η_{μ_1} and η_{μ_2} , with η_1 and η_2 being the so-called *subordination functions*. These functions play a key role in the analytical study of free convolutions [3,4,10].

The functions η_1 and η_2 are unique and can be viewed as η -transforms of certain probability measures on $\mathbb{R}_+$ which are not concentrated at zero. This defines a binary operation $\boxtimes$ on $\mathcal{M}_{\mathbb{R}_+} \setminus \{\delta_0\}$, namely

$$(1.4) \quad \eta_1(z) = \eta_{\mu_2 \boxtimes \mu_1}(z), \quad \text{and} \quad \eta_2(z) = \eta_{\mu_1 \boxtimes \mu_2}(z).$$

The associated convolution $\mu_1 \boxtimes \mu_2$, introduced in this paper, is called the *s-free multiplicative convolution* and it plays the role of the multiplicative analog of the s-free additive convolution $\mu_1 \boxplus \mu_2$ studied in [16].

The subordination formulas (1.3) are related to the so-called *monotone multiplicative convolution* of probability measures on $\mathbb{R}_+$, introduced and studied by Bercovici [6]. This convolution can be defined by the equation

$$(1.5) \quad \eta_{\mu_1 \circ \mu_2}(z) = \eta_{\mu_1}(\eta_{\mu_2}(z))$$

for $z \in \mathbb{C} \setminus \mathbb{R}_+$, where $\mu_1, \mu_2 \in \mathcal{M}_{\mathbb{R}_+}$. Using this convolution and the s-free multiplicative convolution, we can write subordination equations (1.3) in the convolution form

$$(1.6) \quad \mu_1 \boxtimes \mu_2 = \mu_1 \circ (\mu_2 \boxtimes \mu_1) = \mu_2 \circ (\mu_1 \boxtimes \mu_2)$$

for $\mu_1, \mu_2 \in \mathcal{M}_{\mathbb{R}_+} \setminus \{\delta_0\}$.

Further, we show that one can decompose both s-free and free multiplicative convolutions of compactly supported measures on $\mathbb{R}_+$ which are not concentrated at zero in terms of simpler convolutions. For that purpose we introduce another convolution of $\mu_1, \mu_2 \in \mathcal{M}_{\mathbb{R}_+} \setminus \{\delta_0\}$ called the *orthogonal multiplicative convolution*, denoted $\mu_1 \angle \mu_2$. It plays the role of a multiplicative analog of the orthogonal additive convolution on $\mathbb{R}$, introduced and studied in [16]. Using transforms, one can define this convolution by

$$(1.7) \quad \eta_{\mu_1 \angle \mu_2}(z) = \frac{z \eta_{\mu_1}(\eta_{\mu_2}(z))}{\eta_{\mu_2}(z)}$$

for $z \in \mathbb{C} \setminus \mathbb{R}_+$. Using this convolution, we obtain a decomposition of the s-free multiplicative convolution of $\mu_1 \boxtimes \mu_2$ in terms of an infinite sequence of alternating μ_1, μ_2 if these are compactly supported. In view of (1.6), this leads to a decomposition of the free multiplicative convolution.

In a similar way one can treat probability measures on the unit circle $\mathbb{T}$, denoted $\mathcal{M}_{\mathbb{T}}$. In that case, η_μ is the integral over $\mathbb{T}$ and z lies inside the open unit disc $\mathbb{D}$. One defines subordination functions for $\mu_1, \mu_2 \in \mathcal{M}_*$, where $\mathcal{M}_* = \mathcal{M}_{\mathbb{T}} \setminus \{\mu : \mu(X) = 0\}$. Let us also observe that a number of results hold for distributions $\mu_1, \mu_2 \in \Sigma$, where Σ denotes the set functionals $\mu : \mathbb{C}[X] \rightarrow \mathbb{C}$, which send 1 into 1. In that case, the functions η_μ are understood as formal power series.

One of the main points of this paper is that our study of the relations between subordination functions and their decompositions uses operatorial techniques. Namely, we construct bounded operators on Hilbert spaces, which correspond to all compactly supported convolutions which appear in the subordination equations and in the decompositions of s-free convolutions. Our approach is based on the decomposition of the free product of Hilbert spaces $(\mathcal{H}, \xi) = (\mathcal{H}_1, \xi_1) * (\mathcal{H}_2, \xi_2)$, as (two different) orthogonal direct sums

$$(1.8) \quad \mathcal{H} = \bigoplus_{n=1}^{\infty} \mathcal{H}^{(n-1)}(\iota) \oplus \mathcal{H}^{(n)}(\bar{\iota})$$

for each $\iota \in I = \{1, 2\}$, where $\bar{1} = 2$ and $\bar{2} = 1$, with $\mathcal{H}^{(n)}(\iota)$ denoting the subspace spanned by alternating tensor products of order n which do not end with a vector from $\mathcal{H}_\iota^0 := \mathcal{H}_\iota \ominus \mathbb{C}\xi_\iota$, following the original notation of Voiculescu [21] (see also [25]), and we set $\mathcal{H}^{(0)}(\iota) = \mathbb{C}\xi$ for each $\iota \in I$. By $P_\iota(n)$ we denote the canonical projection from $\mathcal{H}$ onto $\mathcal{H}^{(n-1)}(\iota) \oplus \mathcal{H}^{(n)}(\bar{\iota})$ and we set P_ξ to be the projection onto $\mathbb{C}\xi$. Finally, we define the so-called *vacuum state* $\varphi(\cdot) = \langle \cdot, \xi \rangle$ on $\mathcal{B}(\mathcal{H})$.

According to the above decomposition, we can represent bounded free random variables in the form of ‘orthogonal series’

$$(1.9) \quad X_\iota = \sum_{j=1}^{\infty} X_\iota(j),$$

where $\iota \in I$ and $X_\iota(j) = P_\iota(j)X_\iota P_\iota(j)$. These series play a crucial role in our study of subordination for both free additive convolutions [16] and free multiplicative convolutions, studied in this paper. Let us remark that the ‘orthogonal series’ were introduced in a more general context of ‘monotone closed *-algebras’ of operators ‘affiliated’ with unital *-algebras [14]. In that approach, we used the ‘monotone tensor product’ $\bar{\otimes}$ (reminding the von Neumann tensor product) to represent free random variables.

Let μ_1, μ_2 be φ -distributions of bounded random variables X_1 and X_2 , respectively, which are free with respect to φ . In the case of free additive convolutions, crucial was the decomposition of their sum in terms of ‘additive subordination branches’, namely $X_1 + X_2 = B_1 + B_2$, with

$$(1.10) \quad B_\iota = S_\iota^{\text{odd}} + S_\iota^{\text{even}},$$

where

$$(1.11) \quad S_\iota^{\text{odd}} = \sum_{j \text{ odd}} X_\iota(j) \quad \text{and} \quad S_\iota^{\text{even}} = \sum_{j \text{ even}} X_\iota(j),$$

for each $\iota \in I$. Although the subordination branch B_ι is a bounded operator on the ‘free Fock space’ $\mathcal{H}$ for each $\iota \in I$, it acts non-trivially only on its subspace, the ‘s-free Fock space’ $\mathcal{K}_\iota$, defined as

$$(1.12) \quad \mathcal{K}_\iota = \bigoplus_{n \text{ odd}} \mathcal{H}^{(n-1)}(\iota) \oplus \mathcal{H}^{(n)}(\bar{\iota}),$$

and therefore, it is appropriate to view B_ι as an element of $B(\mathcal{K}_\iota)$.

For operatorial subordination in the mutliplicative case, we need to modify the operators S_ι^{odd} , S_ι^{even} and take

$$(1.13) \quad R_\iota^{\text{odd}} = S_\iota^{\text{odd}} \quad \text{and} \quad R_\iota^{\text{even}} = P_\xi + S_\iota^{\text{even}}.$$

Moreover, in order to include all bounded positive random variables, we shall need the operation $\boxtimes$ to be defined for all compactly supported $\mu_1, \mu_2 \in \mathcal{M}_{\mathbb{R}_+}$. For that purpose we set

$$(1.14) \quad \mu \boxtimes \delta_0 = \delta_{\mu(X)} \quad \text{and} \quad \delta_0 \boxtimes \mu = \delta_0$$

for compactly supported $\mu \in \mathcal{M}_{\mathbb{R}_+}$, which turns out natural in our operatorial setting.

Using these notations, and choosing, for simplicity, an existence-type formulation, we can summarize our subordination result for positive operators as follows.

SUBORDINATION FOR POSITIVE OPERATORS. *If X_1 and X_2 are positive, then there exists a decomposition $\sqrt{X_1}X_2\sqrt{X_1} = \sqrt{y}Y\sqrt{y}$, such that the φ -distributions of y and Y are μ_1 and $\mu_2 \boxtimes \mu_1$, respectively, and the pair $(y-1, Y-1)$ is monotone independent w.r.t. φ . Moreover, the φ -distribution of Y agrees with that of $\sqrt{R_1^{\text{even}}R_2^{\text{odd}}}\sqrt{R_1^{\text{even}}}$, and the pair $(R_2^{\text{odd}} - 1_{\mathcal{K}_2}, R_1^{\text{even}} - 1_{\mathcal{K}_2})$ is s-free independent w.r.t. (φ, ψ) , where ψ is the state associated with any unit vector $\zeta \in \mathcal{H}_2^0$.*

Moreover, we further decompose the ‘positive subordination branches’

$$(1.15) \quad \sqrt{R_1^{\text{even}}}R_2^{\text{odd}}\sqrt{R_1^{\text{even}}}$$

using the notion of ‘orthogonal independence’. These decompositions corresponds to decompositions of s-free convolutions in terms of orthogonal convolutions. In a similar way we obtain operatorial subordination results for unitary operators, or even more generally, for bounded operators.

The main examples of operators related to subordination for the free additive convolution, or, more generally, to various notions of independence $\mathcal{I}$, are the adjacency matrices of subgraphs of the corresponding $\mathcal{I}$ -product graphs $\mathcal{G}_1\mathcal{I}\mathcal{G}_2$ (in our study, $\mathcal{I}$ stands for orthogonal, comb, star, s-free or free). In fact, to each independence $\mathcal{I}$, we can associate an additive and a multiplicative convolution,

$$(1.16) \quad \mu_1 +_{\mathcal{I}} \mu_2 \quad \text{and} \quad \mu_1 \times_{\mathcal{I}} \mu_2$$

respectively. Recall that the additive $\mathcal{I}$ -convolution of spectral distributions of rooted graphs corresponds to the addition of ‘monochromatic’ $\mathcal{I}$ -independent adjacency matrices S_ι , $\iota \in I$, and that, in turn, is related to the enumeration of rooted (i.e. root-to-root) walks on $\mathcal{G}_1 \mathcal{I} \mathcal{G}_2$ (for details, see [2] and its references).

In order to find a ‘universal’ multiplicative analog of this theorem, one needs to introduce a new concept of a product of $\mathcal{G}_1$ and $\mathcal{G}_2$ for each $\mathcal{I}$ -independence, which we call the $\mathcal{I}$ loop product and denote $\mathcal{G}_1 \mathcal{I}_\ell \mathcal{G}_2$. Roughly speaking, $\mathcal{G}_1 \mathcal{I}_\ell \mathcal{G}_2$ is obtained by adding colored loops to $\mathcal{G}_1 \mathcal{I} \mathcal{G}_2$ in a suitable way. We assume that the product graph has a ‘natural coloring’, by which we mean that each copy of $\mathcal{G}_\iota$ is colored with color ι , where $\iota \in I$. The procedure of adding loops is equivalent to replacing each S_ι by its ‘unitization’ R_ι obtained from S_ι by adding some projection, in such a way that makes the pair $(R_1 - 1, R_2 - 1)$ $\mathcal{I}$ -independent.

Quite naturally, in order to formulate our result, we shall use the formal power series η_Z corresponding to the variable $Z = R_2 R_1$, namely

$$(1.17) \quad \eta_Z(z) = \sum_{n=1}^{\infty} N_Z(n) z^n,$$

and interpret $N_Z(n)$ as the ‘first return moments’ in the state φ_e associated with the root e . These moments are related to the enumeration of walks of the same class, which we find to be *rooted alternating d-walks* (‘d-walk’ is our abbreviation of ‘double return walks originating with color 1’) counted on different products. Finally, in view of the relation to independence $\mathcal{I}$ just mentioned, these moments agree with the η -moments of the corresponding multiplicative convolutions $\mu_1 \times_{\mathcal{I}} \mu_2$. It is not hard to see that this result can be generalized to the framework of random walks [26].

MULTIPLICATION THEOREM. *Let $\mathcal{G}_1 \mathcal{I}_\ell \mathcal{G}_2$ be naturally colored and let $A(\mathcal{G}_1 \mathcal{I}_\ell \mathcal{G}_2) = R_1 + R_2$ be the decomposition of its adjacency matrix induced by the coloring. Then*

$$(1.17) \quad N_Z(n) = N_{\mu_1 \times_{\mathcal{I}} \mu_2}(n) = |D_{2n}(e)|$$

where $Z = R_2 R_1$ and $D_{2n}(e)$ denotes the set of rooted alternating d-walks on $\mathcal{G}_1 \mathcal{I}_\ell \mathcal{G}_2$ of length $2n$, where $n \in \mathbb{N}$. Moreover, the pair $(R_1 - 1, R_2 - 1)$ is $\mathcal{I}$ -independent.

The paper is organized as follows. In Section 2, we introduce basic notions, including the s-free multiplicative convolution. In Section 3, we introduce and study the concepts of comb- and star loop products of graphs and find relations between rooted alternating d-walks on these graphs and the monotone and boolean multiplicative convolutions, respectively. In Section 4, we introduce and study the orthogonal multiplicative convolution. In Section 5 we define and study the corresponding notion of the orthogonal loop product of rooted graphs. In Section 6 we show, by means of analytical methods, that the definition of the orthogonal multiplicative convolution can be extended to arbitrary measures on $\mathbb{R}_+$ and $\mathbb{T}$. The main operatorial results of the paper are contained in Sections 7 and 8, where we introduce and study operators on the free and s-free Fock spaces which are related to subordination for multiplicative free convolutions and their decompositions. Finally, in Section 9 we find a relation between free and s-free

multiplicative convolutions and the enumeration of rooted alternating d-walks on the free product of graphs and on the s-free loop product of graphs, respectively.

Throughout the whole paper we understand that $I = \{1, 2\}$ and we adopt the notation $\bar{1} = 2$ and $\bar{2} = 1$.

2. PRELIMINARIES

By a *non-commutative probability space* we understand a pair $(\mathcal{A}, \varphi)$, where $\mathcal{A}$ is a unital algebra over $\mathbb{C}$ and φ is a linear functional $\varphi : \mathcal{A} \rightarrow \mathbb{C}$ such that $\varphi(1) = 1$. If $\mathcal{A}$ is a unital $*$ -algebra and φ is positive (called a *state*), then $(\mathcal{A}, \varphi)$ is called a *$*$ -probability space*. If, in addition, $\mathcal{A}$ is a C^* -algebra, then $(\mathcal{A}, \varphi)$ is called a *C^* -probability space*. By the Gelfand-Naimark-Segal theorem, a C^* -probability space can always be realized as a subalgebra of bounded operators on a Hilbert space $\mathcal{H}$ with a distinguished unit vector ξ , for which $\varphi(a) = \langle a\xi, \xi \rangle$ for $a \in \mathcal{A}$.

By a *random variable* we will understand any element a of the considered algebra $\mathcal{A}$. If $\mathcal{A}$ is equipped with an involution, then a random variable a will be called *self-adjoint* if $a = a^*$. The φ -*distribution* of a random variable a is the functional $\mu_a : \mathbb{C}[X] \rightarrow \mathbb{C}$ given by $\mu_a(1) = 1$, $\mu_a(X^n) = \varphi(a^n)$. In particular, if $(\mathcal{A}, \varphi)$ is a C^* -probability space, then the distribution μ_a of a self-adjoint random variable $a \in \mathcal{A}$ extends to a compactly supported probability measure μ on the real line. In that case we will often use the same notation μ for both the distribution of a and the associated compactly supported probability measure.

Various notions of ‘independence’ $\mathcal{I}$ lead to several types of convolutions of distributions (probability measures). If we have two random variables, $X_1 \in \mathcal{A}_1$ and $X_2 \in \mathcal{A}_2$ with φ -distributions μ_1 and μ_2 , respectively, where $\mathcal{A}_1$ and $\mathcal{A}_2$ are $\mathcal{I}$ -independent subalgebras of a noncommutative probability space $(\mathcal{A}, \varphi)$, then the additive convolution of μ and ν associated with $\mathcal{I}$ -independence is the φ -distribution of the sum $X_1 + X_2$. In turn, the multiplicative convolution of μ and ν is the φ -distribution of $X_2 X_1$. In this paper we are interested in the free and s-free multiplicative convolutions associated with free and s-free independence, respectively. In order to decompose them, we shall use the monotone multiplicative convolution introduced by Bercovici [6] and we will introduce the orthogonal multiplicative convolution associated with the notion of ‘orthogonal independence’ [16].

Definition 2.1. Let $\mu_1, \mu_2 \in \mathcal{M}_{\mathbb{R}_+} \setminus \{\delta_0\}$ and let η_1 and η_2 be the associated subordination functions. The unique probability measures $\sigma_1, \sigma_2 \in \mathcal{M}_{\mathbb{R}_+} \setminus \{\delta_0\}$ such that $\eta_1 = \eta_{\sigma_2}$ and $\eta_2 = \eta_{\sigma_1}$ will be called the *s-free multiplicative convolution* of μ_1, μ_2 and μ_2, μ_1 , respectively. We set $\sigma_1 = \mu_1 \boxtimes \mu_2$ and $\sigma_2 = \mu_2 \boxtimes \mu_1$. In a similar way we define the s-free multiplicative convolution of $\mu_1, \mu_2 \in \mathcal{M}_*$.

The s-free multiplicative convolution defines a binary operation $\boxtimes$ on both $\mathcal{M}_{\mathbb{R}_+} \setminus \{\delta_0\}$ and $\mathcal{M}_*$. It can be seen that it is neither commutative nor associative. We will show later that it is related to the concept of ‘freeness with subordination’, or ‘s-free independence’, introduced in [16] (a relation between freeness and monotone independence was also studied in [15]). Moreover, it allows us to rewrite (1.3) in terms of convolutions.

Proposition 2.1. *If $\mu_1, \mu_2 \in \mathcal{M}_{\mathbb{R}_+} \setminus \{\delta_0\}$, then*

$$(2.1) \quad \mu_1 \boxtimes \mu_2 = \mu_1 \cup (\mu_2 \boxtimes \mu_1) = \mu_2 \cup (\mu_1 \boxtimes \mu_2)$$

is the decomposition of the free multiplicative convolution corresponding to the subordination equations (1.3). A similar result holds for $\mu_1, \mu_2 \in \mathcal{M}_*$.

Proof. This fact is an immediate consequence of (1.3), in view of (1.5). ■

By a *rooted set* we understand a pair (X, e) , where X is a countable set and e is a distinguished element of X called *root*. By a *rooted graph* we understand a pair $(\mathcal{G}, e)$, where $\mathcal{G} = (V, E)$ is a non-oriented graph with the set of *vertices* $V = V(\mathcal{G})$, and the set of edges $E = E(\mathcal{G}) \subseteq \{(x, x') : x, x' \in V\}$ and $e \in V$ is a distinguished vertex called the *root*. We identify (x, x') with (x', x) since we consider non-oriented graphs, but when speaking of walks we find it convenient to say that (x, x') *begins* with x and *terminates* with x' . Another reason for using this notation for edges is that of main interest to us are graphs which have *loops*, i.e. edges of the form (x, x) , where $x \in V$. The notion of a rooted graph can be easily generalized to allow multiple edges. Formally, we then obtain a *rooted multigraph*, but we will still use the term ‘rooted graph’, or simply ‘graph’ since all graphs will be considered to have a root. Moreover, we will very often omit the root in our notation and denote by $\mathcal{G}$ the rooted graph $(\mathcal{G}, e)$ if no confusion arises. Thus, in a graph $\mathcal{G}$ we denote by $n(x, x')$ the number of edges connecting x and x' (it may be zero).

For (rooted) graphs we will also use the notation

$$(2.2) \quad V^0 = V \setminus \{e\}.$$

Two vertices $x, x' \in V$ are called *adjacent* if x and x' are connected by an edge, which we denote $x \sim x'$. The *degree* of $x \in V$ is defined by $\kappa(x) = \sum_{x' \sim x} n(x, x')$. A graph is called *locally finite* if $\kappa(x) < \infty$ for every $x \in V$. It is called *uniformly locally finite* if $\sup\{\kappa(x) : x \in V\} < \infty$. All graphs considered in this paper will be connected and uniformly locally finite.

For $x \in V$, let $\delta(x)$ be the indicator function of the one-element set $\{x\}$. Then $\{\delta(x), x \in V\}$ is an orthonormal basis of the Hilbert space $l^2(V)$ of square integrable functions on the set V , with the usual inner product. The *adjacency matrix* $A = A(\mathcal{G})$ of a graph $\mathcal{G}$ is a matrix defined by $A(\mathcal{G})_{x, x'} = n(x, x')$. We identify A with the densely defined symmetric operator on $l^2(V)$ defined by

$$(2.3) \quad A(\mathcal{G})\delta(x) = \sum_{x \sim x'} n(x, x')\delta(x')$$

for $x \in V$. Notice that the sum on the right-hand-side is finite since our graph is assumed to be locally finite. It is known that $A(\mathcal{G})$ is bounded iff $\mathcal{G}$ is uniformly locally finite.

By the *spectral distribution of $A(\mathcal{G})$ in a state ψ* on $l_2(V)$ we understand the measure μ for which

$$(2.4) \quad \psi(A^n) = \int_{\mathbb{R}} x^n \mu(dx), \quad n \in \mathbb{N} \cup \{0\},$$

and by the *spectral distribution of the rooted graph* $(\mathcal{G}, e)$ we understand the spectral distribution of $A(\mathcal{G})$ in the state $\varphi_e(\cdot) = \langle \cdot, \delta(e), \delta(e) \rangle$.

A *walk* from v_0 to v_n on a graph $\mathcal{G}$ is an alternating sequence of vertices and edges of the form

$$(2.5) \quad w = (v_0, \beta_1, v_1, \beta_2, v_2, \dots, v_{n-1}, \beta_n, v_n),$$

with vertices $v_0, v_1, \dots, v_n \in V(\mathcal{G})$ and edges $\beta_1, \beta_2, \dots, \beta_n \in E(\mathcal{G})$, such that β_i is an edge connecting v_{i-1} and v_i . The length of w is given by $l(w) = n$. We allow $v_{i-1} = v_i$, in particular this happens if β_i is a loop.

A *subwalk* of w is a subsequence of w of the form

$$(2.6) \quad w' = (v_i, \beta_i, v_{i+1}, \beta_{i+1}, \dots, \beta_{j-1}, v_j),$$

where $0 \leq i < j \leq n$. In the case when $\mathcal{G}$ does not have multiple edges, we can identify w with the sequence $(v_0, v_1, \dots, v_n)$ with the understanding that $v_{i-1} \sim v_i$ (in this case there is no confusion which edge connecting v_{i-1} and v_i is chosen) for $1 \leq i \leq n$.

The set of walks from v to v' will be denoted $W(v, v')$ and we set $W(v, v) = W(v)$. A walk from $v = v_0$ to $v = v_n$, in which $v_i \neq v$ for all $1 \leq i \leq n-1$ will be called an *f-walk*. Note that we do not require all vertices $v_0, v_1, \dots, v_{n-1}$ in an f-walk to be distinct. The set of f-walks from v to v will be denoted $F(v)$. Subsets of $W(v)$ and $F(v)$ consisting of walks and f-walks of length n , respectively, will be denoted $W_n(v)$ and $F_n(v)$, respectively.

A walk w can also be represented as a sequence of subwalks; in particular, an alternating sequence of edges and subwalks of the form

$$(2.7) \quad w = (w_0, \beta_1, w_1, \beta_2, w_2, \dots, w_{r-1}, \beta_r, w_r),$$

where w_0 is a walk from v_0 to some $v_{i(1)}$, β_1 is an edge connecting $v_{i(1)}$ with $v_{i(1)+1}$, w_1 is a walk from $v_{i(1)+1}$ to some $v_{i(2)}$, β_2 is an edge connecting $v_{i(2)}$ with $v_{i(2)+1}$, and so on, finally, w_r is a walk from $v_{i(r)+1}$ to $v_{i(r+1)} = v_n$. We shall also use similar representations of walks which begin or end with an edge. Representing a walk in terms of edges and subwalks is very convenient when the subwalks $w_1, w_2, \dots, w_r$ are of special type (for instance, are f-walks).

A graph $\mathcal{G}$ whose edges are colored with colors from the set I will be called an *I-edge-colored* graph. In our case, we will always assume that $I = \{1, 2\}$, so we can also use the term ‘2-edge-colored graph’. A walk in such a graph will be called *alternating* if the colors of its edges alternate.

Definition 2.2. Let $\mathcal{G}$ be I -colored. A walk $w \in W(v)$ on $\mathcal{G}$ will be called an *alternating double return walk*, or simply an *alternating d-walk*, if it is alternating, begins with an edge of color 1, and can be represented as a pair (u_1, u_2) of subwalks, such that $u_1, u_2 \in F(v)$. The set of alternating d-walks from v to v will be denoted $D(v)$, and we set $D_m(v) = W_m(v) \cap D(v)$.

Edge-coloring by a two-element set is natural in the case of many products of rooted graphs. In particular, the products of rooted graphs of type $\mathcal{G}_1 \mathcal{I} \mathcal{G}_2$ have the property that every edge of the product graph belongs either to a copy of $\mathcal{G}_1$, or to a copy of $\mathcal{G}_2$. Thus we can color the edges of all copies of $\mathcal{G}_\iota$ with color ι , where $\iota \in I$, in which case

we will say that $\mathcal{G}_1 \mathcal{I} \mathcal{G}_2$ is *naturally colored*. In turn, we will say that the product $\mathcal{G}_1 \mathcal{I}_\ell \mathcal{G}_2$ is *naturally colored* if its coloring is inherited from $\mathcal{G}_1 \mathcal{I} \mathcal{G}_2$. Let us also remark that our condition that an alternating walk should begin with an edge of color 1 (and thus end with an edge of color 2) is caused by the fact that we want to identify the given walk w and its ‘reverse’ obtained by reversing the order in (2.12).

We end this Section with elementary formulas for the ‘first return moments’ of a random variable X , by which we understand the coefficients of the formal power series η_X associated with the distribution of X . If X is a random variable in a non-commutative probability space $(\mathcal{A}, \varphi)$, then the ‘moment generating function’ and the ‘first return moment generating function’, respectively, associated with the φ -distribution μ of X , are given by formal power series

$$(2.8) \quad \psi_X(z) = \sum_{n=1}^{\infty} \mu(X^n)(n) z^n, \quad \eta_X(z) = \frac{\psi_X(z)}{1 + \psi_X(z)} = \sum_{n=1}^{\infty} N_X(n) z^n,$$

where the numbers $N_X(n)$, $n \in \mathbb{N}$, will sometimes be called ‘first return moments of X ’.

Below we give a convenient algebraic formula for these ‘first return moments’. For that purpose, let us take the extension of $(\mathcal{A}, \varphi)$ to a larger noncommutative probability space, namely, the free product with identified units $\mathcal{A} * \mathbb{C}[P]$, where $P^2 = P$, with the state given by the linear extension of

$$(2.9) \quad \varphi(P) = 1, \quad \varphi(w_1 P w_2) = \varphi(w_1) \varphi(w_2)$$

for any $w_1, w_2 \in \mathcal{A} * \mathbb{C}[P]$, where, slightly abusing notation, we also denote the new state by φ [13]. In particular, in the C^* -algebra context, P can be interpreted as the projection onto $\mathbb{C}\xi$, where ξ is the cyclic unit vector of the GNS triple associated with $(\mathcal{A}, \varphi)$.

Proposition 2.2. *Let X be a random variable in a noncommutative probability space $(\mathcal{A}, \varphi)$. Then*

$$(2.10) \quad N_X(n) = \varphi(X(P^\perp X)^{n-1})$$

for $n \in \mathbb{N}$, where $P^\perp = 1 - P$.

Proof. This is a straightforward consequence of (1.2). ■

Finally, let us prove an elementary fact about a relation between the ‘first return moments’ of products of ‘monochromatic’ adjacency matrices in certain I -colored graphs and the cardinalities of the sets $D_{2n}(e)$.

Proposition 2.3. *Let $(\mathcal{G}, e)$ be an I -colored uniformly locally finite graph, and let $A(\mathcal{G}) = A_1 + A_2$ be the decomposition of $A(\mathcal{G})$ induced by the coloring. If the set of rooted alternating f -walks of even lengths is empty, then*

$$(2.11) \quad N_Z(n) = |D_{2n}(e)|,$$

where $Z = A_2 A_1$ and the numbers $N_Z(n)$ correspond to the state $\varphi_e(\cdot) = \langle \cdot, \delta(e), \delta(e) \rangle$.

Proof. If we take $Z = A_2 A_1$ in Proposition 2.2, then $N_Z(n)$ is equal to the number of rooted alternating walks of length $2n$, which begin with an edge of color 1 and such that intermediate returns to the root can occur only after odd numbers of steps. If there

are no rooted alternating f-walks of even length, then the first return to the root must occur after an odd number of steps and thus the second return to the root must occur after an even number of steps. Therefore, if we consider rooted alternating walks which have an even number of steps and such that intermediate returns occur only after an odd number of steps, these are d-walks. This proves our assertion. ■

Let us note that all graph products considered in this paper satisfy the assumptions of Proposition 2.3, and therefore, walk-counting is always reduced to rooted alternating d-walks.

3. COMB AND STAR LOOP PRODUCTS OF GRAPHS

We begin with recalling the notions of the additive [18] and multiplicative [6] monotone convolutions and show that the η -moments of the latter are related to the enumeration of alternating d-walks on a new version of the comb product of graphs called the ‘comb loop product’.

Definition 3.1. Two subalgebras $\mathcal{A}_1, \mathcal{A}_2$ of a unital algebra $\mathcal{A}$ are *monotone independent* with respect to a normalized linear functional φ on $\mathcal{A}$ if

$$\begin{aligned} \varphi(w_1 a_1 b) &= \varphi(b) \varphi(w_1 a_1) \quad \text{and} \quad \varphi(b a_2 w_2) = \varphi(b) \varphi(a_2 w_2), \\ \varphi(w_1 a_1 b a_2 w_2) &= \varphi(b) \varphi(w_1 a_1 a_2 w_2), \end{aligned}$$

whenever $a_1, a_2 \in \mathcal{A}_1$, $b \in \mathcal{A}_2$ and w_1, w_2 are arbitrary elements of the unital algebra $\text{alg}(\mathcal{A}_1, \mathcal{A}_2)$ generated by $\mathcal{A}_1$ and $\mathcal{A}_2$. In particular, we will say that the pair (a, b) of elements of $\mathcal{A}$ is monotone independent w.r.t. φ if the algebras generated by these elements are monotone independent.

Suppose φ -distributions of random variables a_1 and a_2 are μ_1 and μ_2 , respectively. If the pair (a_1, a_2) is monotone independent w.r.t. φ , then the φ -distribution of the sum $a_1 + a_2$ is the monotone additive convolution $\mu_1 \triangleright \mu_2$. In turn, if the pair $(a_1 - 1, a_2 - 1)$ is monotone independent w.r.t. φ , then the φ -distribution of the product $a_2 a_1$ is the *monotone multiplicative convolution* $\mu_1 \cup \mu_2$. The corresponding formal power series satisfy (1.5). If $\mu_1, \mu_2 \in \mathcal{M}_{\mathbb{R}_+}$ or $\mu_1, \mu_2 \in \mathcal{M}_{\mathbb{T}}$, then (1.5) is understood in terms of analytic self-maps of $\mathbb{C} \setminus \mathbb{R}_+$ or $\mathbb{T}$, respectively. For details, see [19] (additive case) and [6] (multiplicative case).

It is known that the monotone independence can be associated with the comb product of graphs [1]. Let us recall the definition of the comb product and follow up with the corresponding loop product.

Definition 3.2. The *comb product* of rooted graphs $(\mathcal{G}_1, e_1)$ and $(\mathcal{G}_2, e_2)$ is the rooted graph $(\mathcal{G}_1 \triangleright \mathcal{G}_2, e)$ obtained by attaching a copy of $\mathcal{G}_2$ by its root e_2 to each vertex of $\mathcal{G}_1$, where we denote by e the vertex obtained by identifying e_1 and e_2 . If no confusion arises, we denote the comb product by $\mathcal{G}_1 \triangleright \mathcal{G}_2$. If we identify its set of vertices with $V_1 \times V_2$, then its root is identified with $e_1 \times e_2$.

Definition 3.3. Suppose the edges of the comb product $(\mathcal{G}_1 \triangleright \mathcal{G}_2, e)$ are naturally colored. The *comb loop product* of rooted graphs $(\mathcal{G}_1, e_1)$ and $(\mathcal{G}_2, e_2)$ is the rooted graph $(\mathcal{G}_1 \triangleright_\ell \mathcal{G}_2, e)$ obtained from $(\mathcal{G}_1 \triangleright \mathcal{G}_2, e)$ by attaching a loop of color 1 to each vertex but the root of each copy of $\mathcal{G}_2$.

Let us justify the above definition. We follow the observation made by Bercovici [6] that in order to introduce a multiplicative convolution associated with monotone independence, one needs to ‘unitize’ the usual monotone independent variables. It has been shown in [5] that one of the possible choices is to take variables of the form

$$(3.1) \quad R_1 = a_1 \otimes P_2 + 1_1 \otimes P_2^\perp \quad \text{and} \quad R_2 = 1_1 \otimes a_2,$$

where $a_\iota, 1_\iota \in B(\mathcal{H}_\iota)$, with $(\mathcal{H}_\iota, \xi_\iota)$, being Hilbert spaces with distinguished unit vectors and identity operators 1_ι , and $\iota \in I$. Then the moments of the product $R_2 R_1$ in the state associated with the vector $\xi_1 \otimes \xi_2$ agree with the moments of $\mu_1 \cup \mu_2$.

If a_ι is taken to be the adjacency matrix of a (uniformly locally finite) graph $\mathcal{G}_\iota$, $\iota \in I$, then the term $1_1 \otimes P_2^\perp$ corresponds to glueing a loop of color 1 to each vertex but the root of each copy of $\mathcal{G}_2$. Thus, R_1 is obtained from the usual comb product adjacency matrix S_1 by adding a projection L_1 , whereas $R_2 = S_2$, where

$$(3.2) \quad S_1 = a_1 \otimes P_2, \quad L_1 = 1_1 \otimes P_2^\perp$$

and R_1, R_2 are the adjacency matrices of ‘monochromatic’, 1 and 2-colored subgraphs, respectively. This realization can be generalized to arbitrary random variables in non-commutative probability spaces, using the extensions of functionals given by (2.9).

Remark 3.1. It is not difficult to show that the comb loop product of graphs $\mathcal{G}_1$ and $\mathcal{G}_2$ is isomorphic to the usual comb product of larger graphs $\tilde{\mathcal{G}}_1$ and $\tilde{\mathcal{G}}_2$. However, in order to keep the same coloring of the product graph (which is needed for counting alternating walks) one has to use two colors for $\tilde{\mathcal{G}}_2$. Namely, $\tilde{\mathcal{G}}_2$ is obtained from $\mathcal{G}_2$ by glueing a loop of color 1 to each vertex but the root of $\mathcal{G}_2$. Then we indeed have $\mathcal{G}_1 \triangleright_\ell \mathcal{G}_2 = \tilde{\mathcal{G}}_1 \triangleright \tilde{\mathcal{G}}_2$, but usefulness of this relation seems to be limited (roughly speaking, we have a simpler product but a more complicated coloring).

Example 3.1. Let us consider the example of the comb loop product of graphs given in Figure 1. We choose both $\mathcal{G}_1$ and $\mathcal{G}_2$ to have a loop at the root (to distinguish the loops, we draw loops of color 1 smaller than loops of color 2, which is helpful in the enumeration of alternating walks). Then $\mathcal{G}_1 \triangleright_\ell \mathcal{G}_2$ has loops of two types: loops whose origin can be traced back to graphs $\mathcal{G}_1$ and $\mathcal{G}_2$ (these are all loops which are at the glueing points), or are added to the usual comb product in the process of forming the comb loop product. Denote by $(v, v)_\iota$ the loop from v to v of color ι . The following walk is a rooted alternating d-walk:

$$(\beta_1, w_1, \beta_2, w_2, \beta_3, w_3, \beta_4, w_4) \in D_8(e),$$

where the sequence of edges of color 1,

$$(\beta_1, \beta_2, \beta_3, \beta_4) = ((e, x), (x, x'), (x', x), (x, e))$$

forms a rooted f-walk of color 1, which is interlaced with the sequence (w_1, w_2, w_3, w_4) of loops of color 2: $w_1 = w_3 = (x, x)_2 \in F_1(x)$, $w_2 = (x', x')_2 \in F_1(x')$ and $w_4 = (e, e)_2 \in$




FIGURE 1. An example of the comb loop product

$F_1(e)$. Note that the first return to e occurs at the end of β_4 , whereas the second return occurs as the end of w_4 . When any of the loops, w_i , $1 \leq i \leq 4$, is replaced by an arbitrary alternating f-walk from the corresponding vertex to itself which begins and ends with an edge of color 2, we still get a rooted alternating d-walk. The simplest examples of a rooted alternating d-walk are of course given by: $(\beta_0, w_0) \in D_2(e)$, where $\beta_0 = (e, e)_1$, $w_0 = (e, e)_2$ and $(\beta_1, w_1, \beta_4, w_4) \in D_4(e)$.

Theorem 3.1. *The Multiplication Theorem holds for monotone independence, the associated loop product $\mathcal{G}_1 \triangleright_\ell \mathcal{G}_2$, and the multiplicative convolution $\mu_1 \circ \mu_2$.*

Proof. From Definition 3.3 and (3.1) it follows that

$$Z = A(\mathcal{G}_1 \triangleright_\ell \mathcal{G}_2) = R_1 + R_2,$$

where the pair $(R_1 - 1, R_2 - 1)$ is monotone independent w.r.t. φ_e , the state associated with e and R_ι , $\iota \in I$, is the adjacency matrix of the ι -colored subgraph. Moreover, by Proposition 2.2, $N_Z(n)$ is the number of alternating d-walks on $\mathcal{G}_1 \triangleright_\ell \mathcal{G}_2$ of length $2n$. This proves the first equation in (1.13) for monotone independence. Now, from equation (1.5), we obtain the combinatorial formula

$$N_{\mu_1 \circ \mu_2}(n) = \sum_{r=1}^n N_{\mu_1}(r) \sum_{k_1+k_2+\dots+k_r=n} N_{\mu_2}(k_1)N_{\mu_2}(k_2)\dots N_{\mu_2}(k_r),$$

where it is tacitly assumed that the indices $k_1, k_2, \dots, k_r$ are positive integers and numbers $N_\mu(n)$ are coefficients of η_μ treated as formal power series

$$\eta_\mu(z) = \sum_{n=1}^{\infty} N_\mu(n)z^n.$$

The above combinatorial formula allows us to find a correspondence between these η -moments and rooted alternating d-walks on $\mathcal{G}_1 \triangleright_\ell \mathcal{G}_2$. Let us observe that $N_{\mu_1}(r)$ is the number of rooted f-walks of length r on $\mathcal{G}_1$ and $N_{\mu_2}(k)$ is the number of rooted f-walks of length k on $\mathcal{G}_2$. On the other hand, recall that in the comb loop product $\mathcal{G}_1 \triangleright_\ell \mathcal{G}_2$ there is only one copy of $\mathcal{G}_1$ (with e_1 identified with the root e of the product graph), with a copy of $\mathcal{G}_2$ attached by its root to every vertex x of $\mathcal{G}_1$ (the vertex x becomes then the only common vertex of this copy of $\mathcal{G}_2$ and the original copy of $\mathcal{G}_1$). In addition, loops of color 1 are glued to all vertices but the roots of these copies of $\mathcal{G}_2$. Therefore, each rooted alternating d-walk on $\mathcal{G}_1 \triangleright_\ell \mathcal{G}_2$ consists of a sequence of edges of $\mathcal{G}_1$ which forms an f-walk of color 1, namely $c = (v_0, v_1, v_2, \dots, v_{r-1}, v_r) \in F(e)$, and alternating f-walks $c_i \in F(v_i)$, $i = 1, \dots, r$, with the first edge of color 2, attached to every vertex

of c . Note that the number of alternating f-walks from v_i to v_i with the first edge of color 2 is equal to the number of walks from e_2 to e_2 in graph $\mathcal{G}_2$ – the only difference is that if the latter has length k_i , the former has length $2k_i - 1$ since each edge must be followed by a loop of color 2. Thus, the contribution from each product of the form

$$N_{\mu_1}(r)N_{\mu_2}(k_1)N_{\mu_2}(k_2)\dots N_{\mu_2}(k_r)$$

to the RHS of the formula for $N_{\mu_1 \circ \mu_2}(n)$ is equal to the number of all such f-walks c of color 1 which have r edges, interlaced with r alternating f-walks of lengths $2k_1 - 1, 2k_2 - 1, \dots, 2k_r - 1$. The summation over $k_1 + k_2 + \dots + k_r = n$ indicates that all alternating f-walks involved must have $2n - r$ edges together. The summation over $1 \leq r \leq n$ gives exactly the cardinality of $D_{2n}(e)$, which finishes the proof. $\blacksquare$

Let us observe here that (3.1) can be generalized to arbitrary random variables in noncommutative probability spaces with distributions $\mu_1, \mu_2 \in \Sigma$ if one treats P_1 and P_2 as idempotents and uses extensions of states φ_1, φ_2 given by (2.9). Then, the combinatorial formula in the proof of Theorem 3.1 remains valid for $\mu_1, \mu_2 \in \Sigma$.

Example 3.2. If $\mu_1, \mu_2 \in \Sigma$, the lowest order moments of $N_{\mu_1 \circ \mu_2}$ are given by

$$\begin{aligned} N_{\mu_1 \circ \mu_2}(1) &= N_{\mu_1}(1)N_{\mu_2}(1), \\ N_{\mu_1 \circ \mu_2}(2) &= N_{\mu_1}(2)N_{\mu_2}^2(1) + N_{\mu_1}(1)N_{\mu_2}(2), \\ N_{\mu_1 \circ \mu_2}(3) &= N_{\mu_1}(3)N_{\mu_2}^3(1) + 2N_{\mu_1}(2)N_{\mu_2}(2)N_{\mu_2}(1) + N_{\mu_1}(1)N_{\mu_2}(3), \\ N_{\mu_1 \circ \mu_2}(4) &= N_{\mu_1}(4)N_{\mu_2}^4(1) + 3N_{\mu_1}(3)N_{\mu_2}(2)N_{\mu_2}^2(1) + N_{\mu_1}(2)N_{\mu_2}^2(2) \\ &\quad + 2N_{\mu_1}(2)N_{\mu_2}(3)N_{\mu_2}(1) + N_{\mu_1}(1)N_{\mu_2}(4). \end{aligned}$$

Let us apply these formulas to the graph $\mathcal{G}_1 \supseteq_\ell \mathcal{G}_2$ in Fig.1. Let μ_1 and μ_2 be the spectral distributions associated with $\mathcal{G}_1$ and $\mathcal{G}_2$, respectively. By counting f-walks on $\mathcal{G}_1$ and $\mathcal{G}_2$, we easily get $N_{\mu_1}(1) = N_{\mu_1}(2) = N_{\mu_1}(4) = 1$, $N_{\mu_1}(3) = 0$ and $N_{\mu_2}(1) = N_{\mu_2}(2) = 1$, $N_{\mu_2}(3) = 0$, $N_{\mu_2}(4) = 2$. In turn, using the above formulas and Theorem 3.1, we get $D_2(e) = 1$, $D_4(e) = D_6(e) = 2$ and $D_8(e) = 4$, which can also be verified directly by counting comb walks on $\mathcal{G}_1 \supseteq_\ell \mathcal{G}_2$.

For completeness, let us also briefly discuss the case of the star product of graphs $\mathcal{G}_1 \star \mathcal{G}_2$ and find its relation to the boolean multiplicative convolution.

Definition 3.4. The *star product* of rooted graphs $(\mathcal{G}_1, e_1)$ and $(\mathcal{G}_2, e_2)$ is the rooted graph $(\mathcal{G}_1 \star \mathcal{G}_2, e)$ obtained by glueing $\mathcal{G}_1$ and $\mathcal{G}_2$ at their roots and taking this common root to be the root of the product. The *star loop product* $(\mathcal{G}_1 \star_\ell \mathcal{G}_2, e)$ is obtained from naturally colored $(\mathcal{G}_1 \star \mathcal{G}_2, e)$ by adding loops of color ι to every vertex but the root of $\mathcal{G}_\iota$, where $\iota \in I$.

Recall here the definition of the boolean multiplicative convolution of distributions $\mu_1, \mu_2 \in \Sigma$ [5]. Treating $\eta_\mu(z)$ as a formal power series, we define

$$(3.3) \quad \rho_\mu(z) = \frac{\eta_\mu(z)}{z},$$




FIGURE 2. An example of the star loop product

which allows us to define the boolean multiplicative convolution of $\mu_1, \mu_2 \in \Sigma$ by the formula

$$(3.4) \quad \rho_{\mu_1 \boxtimes \mu_2}(z) = \rho_{\mu_1}(z) \rho_{\mu_2}(z).$$

This formula defines $\boxtimes$ as a binary operation on $\mathcal{M}_{\mathbb{T}}$ [11]. However, if $\mu_1, \mu_2 \in \mathcal{M}_{\mathbb{R}_+}$, then the RHS does not always give the ρ -transform of a probability measure on $\mathbb{R}_+$ [5].

Nevertheless, the well-known relation [8,24]

$$(3.5) \quad \rho_{\mu_1 \boxtimes \mu_2}(z) = \rho_1(z) \rho_2(z)$$

where $\rho_{\iota}(z) = \eta_{\iota}(z)/z$, $\iota \in I$, can still be written in terms of convolutions as

$$(3.6) \quad \mu_1 \boxtimes \mu_2 = (\mu_1 \boxdot \mu_2) \boxtimes (\mu_2 \boxdot \mu_1).$$

which was our motivation to use the symbol $\boxdot$ for the s-free convolution (an analogous formula holds for $\mu_1, \mu_2 \in \mathcal{M}_*$).

If a_{ι} is the adjacency matrix of a graph $\mathcal{G}_{\iota}$, where $\iota \in I$, then, using a similar reasoning as in the case of the comb loop product, we introduce operators of the form

$$\begin{aligned} R_1 &= a_1 \otimes P_2 + 1 \otimes P_2^{\perp}, \\ R_2 &= P_1 \otimes a_2 + P_1^{\perp} \otimes 1_2, \end{aligned}$$

which can be viewed as the adjacency matrices of the 1- and 2-colored subgraphs of the product graph $(\mathcal{G}_1 \star_{\ell} \mathcal{G}_2, e)$. Moreover, these are exactly the operators which give the moments of the boolean multiplicative convolution of spectral distributions of $\mathcal{G}_1$ and $\mathcal{G}_2$ since the pair $(R_1 - 1, R_2 - 1)$ is boolean independent w.r.t. φ_e . An example of a star loop product is given in Fig.2.

Let us also note that formulas (3.3)-(3.4) can be used if a_1, a_2 are random variables in arbitrary noncommutative probability spaces – this is obtained if one uses extended states (2.9). This gives a realization of the boolean multiplicative convolution of arbitrary distributions $\mu_1, \mu_2 \in \Sigma$.

Proposition 3.1. *The Multiplication Theorem holds for boolean independence, the associated loop product $\mathcal{G}_1 \star_{\ell} \mathcal{G}_2$, and the multiplicative convolution $\mu_1 \boxtimes \mu_2$.*

Proof. The proof is similar to that of Theorem 3.1 and is based on the easy combinatorial formula

$$N_{\mu_1 \boxtimes \mu_2}(n) = \sum_{j+k=n+1} N_{\mu_1}(j) N_{\mu_2}(k),$$

where the sum runs over positive indices j, k , obtained from (1.2) and (3.4). Details are left to the reader. ■

Example 3.3. If $\mu_1, \mu_2 \in \Sigma$, the lowest order ‘first return moments’ $N_{\mu_1 \boxtimes \mu_2}(n)$ are given by

$$\begin{aligned} N_{\mu_1 \boxtimes \mu_2}(1) &= N_{\mu_1}(1)N_{\mu_2}(1), \\ N_{\mu_1 \boxtimes \mu_2}(2) &= N_{\mu_1}(1)N_{\mu_2}(2) + N_{\mu_1}(2)N_{\mu_2}(1), \\ N_{\mu_1 \boxtimes \mu_2}(3) &= N_{\mu_1}(1)N_{\mu_2}(3) + N_{\mu_1}(2)N_{\mu_2}(2) + N_{\mu_1}(3)N_{\mu_2}(1), \\ N_{\mu_1 \boxtimes \mu_2}(4) &= N_{\mu_1}(1)N_{\mu_2}(4) + N_{\mu_1}(2)N_{\mu_2}(3) + N_{\mu_1}(3)N_{\mu_2}(2) + N_{\mu_1}(4)N_{\mu_2}(1). \end{aligned}$$

These formulas can be used to count rooted alternating d-walks on the product graph given in Fig. 2. Using Proposition 3.2 and the values of $N_{\mu_1}(j)$ and $N_{\mu_2}(k)$, given in Example 3.1, we get $D_2(e) = D_4(e) = D_6(e) = 1$ and $D_8(e) = 3$, which can be also verified by direct computations.

4. ORTHOGONAL MULTIPLICATIVE CONVOLUTION

For given positive bounded random variables X_1 and X_2 with distributions μ_1 and μ_2 , we want to find a positive random variable Z on some Hilbert space whose distribution is given by the multiplicative analog of the orthogonal convolution, denoted $\mu_1 \angle \mu_2$.

Let us first recall the notion of orthogonal independence [16].

Definition 4.1. Let $(\mathcal{A}, \varphi, \psi)$ be a unital algebra with a pair of linear normalized functionals and let $\mathcal{A}_1$ and $\mathcal{A}_2$ be non-unital subalgebras of $\mathcal{A}$. We say that $\mathcal{A}_2$ is *orthogonal* to $\mathcal{A}_1$ with respect to (φ, ψ) if

- (i) $\varphi(bw_2) = \varphi(w_1b) = 0$,
- (ii) $\varphi(w_1a_1ba_2w_2) = \psi(b)(\varphi(w_1a_1a_2w_2) - \varphi(w_1a_1)\varphi(a_2w_2))$,

for any $a_1, a_2 \in \mathcal{A}_1$, $b \in \mathcal{A}_2$, and any elements w_1, w_2 of the unital algebra $\text{alg}(\mathcal{A}_1, \mathcal{A}_2)$ generated by $\mathcal{A}_1$ and $\mathcal{A}_2$. We say that the pair (a, b) of elements of $\mathcal{A}$ is *orthogonal* with respect to (φ, ψ) if the algebra generated by $b \in \mathcal{A}$ is orthogonal to the algebra generated by $a \in \mathcal{A}$.

Definition 4.2. Let $(\mathcal{A}, \varphi, \psi)$ be a unital algebra with a pair of normalized linear functionals and let $\mathcal{A}_1, \mathcal{A}_2$ be a pair of (in general, non-unital) subalgebras of $\mathcal{A}$, such that $\mathcal{A}_2$ is orthogonal to $\mathcal{A}_1$ w.r.t. (φ, ψ) . Let $a_1, a_2 \in \mathcal{A}$ be random variables with φ -distribution μ_1 and ψ -distribution μ_2 , respectively. The *orthogonal multiplicative convolution* of μ_1 and μ_2 , denoted $\mu_1 \angle \mu_2$, is the distribution of a_2a_1 , where $(a_1 - 1, a_2 - 1)$ is orthogonal w.r.t. (φ, ψ) .

In order to find a Hilbert space realization of the orthogonal multiplicative convolution of compactly supported probability measures, recall the definition of the orthogonal product of two Hilbert spaces with distinguished unit vectors $(\mathcal{H}_1, \xi_1)$ and $(\mathcal{H}_2, \xi_2)$. Thus, the *orthogonal product* of $(\mathcal{H}_1, \xi_1)$ and $(\mathcal{H}_2, \xi_2)$ is the pair $(\mathcal{H}, \xi)$, where

$$(4.1) \quad \mathcal{H} = \mathbb{C}\xi \oplus \mathcal{H}_1^0 \oplus (\mathcal{H}_2^0 \otimes \mathcal{H}_1^0),$$

with $\mathcal{H}_i^0 = \mathcal{H}_i \ominus \mathbb{C}\xi_i$ and where ξ is a unit vector. We denote it by $(\mathcal{H}_1, \xi_1) \vdash (\mathcal{H}_2, \xi_2)$, or simply $\mathcal{H}_1 \vdash \mathcal{H}_2$, if no confusion arises. It has been shown in [16] that $\mathcal{H}$ gives a

Hilbert space realization of orthogonal random variables. For that purpose we defined an isometry $U : \mathcal{H} \rightarrow \mathcal{H}_1 \otimes \mathcal{H}_2$ given by

$$(4.2) \quad U(\xi) = \xi_1 \otimes \xi_2, \quad U(h_1) = h_1 \otimes \xi_2, \quad U(h_2 \otimes h_1) = h_1 \otimes h_2$$

for any $h_1 \in \mathcal{H}_1^0$ and $h_2 \in \mathcal{H}_2^0$. Using U , we define faithful non-unital $*$ -homomorphisms $\tau_i : \mathcal{B}(\mathcal{H}_i) \rightarrow \mathcal{B}(\mathcal{H})$ by

$$(4.3) \quad \tau_1(a_1) = U^*(a_1 \otimes P_2)U, \quad \tau_2(a_2) = U^*(P_1^\perp \otimes a_2)U,$$

where P_1, P_2 are the projections onto $\mathbb{C}\xi_1$ and $\mathbb{C}\xi_2$, respectively. Then the pair $(\tau_1(a_1), \tau_2(a_2))$ is orthogonal w.r.t. (φ, ψ) , where φ and ψ are states on $\mathcal{B}(\mathcal{H})$ associated with the unit vector ξ and any unit vector $\zeta \in \mathcal{H}_1^0$, respectively. For details, see [16].

For simplicity, we will identify vectors from $\mathcal{H}$ with their images in $\mathcal{H}_1 \otimes \mathcal{H}_2$ under the isometry U as well as bounded operators a, b with $\tau_1(a_1), \tau_2(a_2)$, respectively, and we will carry out computations in $\mathcal{H}_1 \otimes \mathcal{H}_2$.

Proposition 4.1. *Let $a_1 \in B(\mathcal{H}_1)$ have φ_1 -distribution μ_1 and $a_2 \in B(\mathcal{H}_2)$ have φ_2 -distribution μ_2 . Then there exist a Hilbert space $\mathcal{H}$ with unit vectors ξ and ζ , and random variables $R_1, R_2 \in B(\mathcal{H})$ such that R_1 has φ -distribution μ_1 , R_2 has ψ -distribution μ_2 and $R_2 R_1$ has φ -distribution $\mu_1 \angle \mu_2$, where φ and ψ are states associated with ξ and ζ , respectively.*

Proof. Let $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2$ with the canonical unit vector $\xi = \xi_1 \otimes \xi_2$ and let $\zeta = \zeta_1 \otimes \xi_2$, where ζ_1 is an arbitrary unit vector from $\mathcal{H}_1^0$. Define

$$\begin{aligned} R_1 &= a_1 \otimes P_2 + 1_1 \otimes P_2^\perp, \\ R_2 &= P_1^\perp \otimes a_2 + P_1 \otimes 1_2, \end{aligned}$$

where $P_i^\perp = 1_i - P_i$ and P_i is the canonical projection onto $\mathbb{C}\xi_i$ and 1_i denotes the identity in $B(\mathcal{H}_i)$, $i = 1, 2$. Therefore, denoting $1 = 1_1 \otimes 1_2$, we get

$$\begin{aligned} R_1 - 1 &= (a_1 - 1_1) \otimes P_2, \\ R_2 - 1 &= P_1^\perp \otimes (a_2 - 1_2), \end{aligned}$$

which, due to the appropriate tensor product form, implies that the pair $(R_1 - 1, R_2 - 1)$ is orthogonal w.r.t. (φ, ψ) , where φ is the state associated with ξ and ψ is the state associated with ζ . Moreover, elementary calculations show that the φ -distribution of R_1 is μ_1 and that the ψ -distribution of R_2 is μ_2 , which completes the proof. $\blacksquare$

Corollary 4.1. *Let μ_1, μ_2 be compactly supported probability measures on $\mathbb{R}_+$. Then there exist bounded positive random variables R_1, R_2 on some Hilbert space $\mathcal{H}$ and unit vectors $\xi, \zeta \in \mathcal{H}$, such that R_1 has φ -distribution μ_1 , R_2 has ψ -distribution μ_2 and $\sqrt{R_2} R_1 \sqrt{R_2}$ has φ -distribution $\mu_1 \angle \mu_2$.*

Proof. There exist positive operators $x_1 \in B(L^2(\mathbb{R}_+, \mu_1))$ and $x_2 \in B(L^2(\mathbb{R}_+, \mu_2))$ (standard multiplication operators) which have distributions μ_1 and μ_2 , respectively, with respect to the states φ_1 and φ_2 associated with the constant functions equal to

one, h_1 and h_2 , respectively. We can then use the framework of Proposition 4.1 to get self-adjoint bounded random variables on $\mathcal{H} = L^2(\mathbb{R}, \mu_1) \otimes L^2(\mathbb{R}, \mu_2)$ of the form

$$\begin{aligned} R'_1 &= (x_1 - 1_1) \otimes P_2, \\ R'_2 &= P_1^\perp \otimes (x_2 - 1_2), \end{aligned}$$

which are orthogonal w.r.t. (φ, ψ) , where φ is associated with vector $\xi = h_1 \otimes h_2$ and ψ is associated with $\zeta = f \otimes h_2$, where f is any function such that $\int_{\mathbb{R}} f(x) d\mu_1(x) = 0$ and $\int_{\mathbb{R}} f^2(x) d\mu_1(x) = 1$. Now, let $R_1 = R'_1 + 1$ and $R_2 = R'_2 + 1$. It follows from the proof of Proposition 4.1 that R_1 and R_2 have distributions μ_1 and μ_2 , respectively. Moreover, both are positive operators and thus, in particular, $\sqrt{R_2}$ exists. In fact,

$$\sqrt{R_2} = P_1^\perp \otimes \sqrt{a_2} + P_1 \otimes 1_2,$$

and $\sqrt{R_2} R_1 \sqrt{R_2}$ is a bounded positive operator on $\mathcal{H}$. Moreover, $\sqrt{R_2} R_1 \sqrt{R_2}$ has the same φ -distribution as $R_1 R_2$, which is obtained from the following calculation:

$$\begin{aligned} \varphi((\sqrt{R_2} R_1 \sqrt{R_2})^n) &= \varphi(\sqrt{R_2} (R_1 R_2)^{n-1} R_1 \sqrt{R_2}) \\ &= (\varphi_1 \otimes \varphi_2)((\sqrt{P_1} \otimes 1_2)(R_1 R_2)^{n-1} R_1 (P_1 \otimes 1_2)) \\ &= (\varphi_1 \otimes \varphi_2)((R_1 R_2)^{n-1} R_1 (P_1 \otimes 1_2)) \\ &= \varphi((R_1 R_2)^n). \end{aligned}$$

By using the adjoints, we can interchange the order of R_1 and R_2 under the symbol φ . Therefore, the probability distribution of $\sqrt{R_2} R_1 \sqrt{R_2}$ extends to the compactly supported measure $\mu_1 \angle \mu_2$ on $\mathbb{R}_+$. $\blacksquare$

We will now compute the φ -distribution of the product $Z := R_2 R_1$, where R_1 and R_2 are given by Proposition 4.1, and derive a formula for η_Z . In this context, let us observe that one can generalize Proposition 4.1 to arbitrary random variables in noncommutative probability spaces since formulas for R_1 and R_2 remain valid provided one takes extensions of states given by (2.9) and uses Proposition 2.2. Therefore, further developments in this Section will hold for arbitrary random variables in noncommutative probability spaces. This corresponds to taking distributions $\mu \in \Sigma$ and the corresponding formal power series η_μ and ρ_μ .

In order to compute the φ -distribution of Z , let us decompose Z as

$$Z = Z_1 + Z_2 + Z_3 + Z_4,$$

where

$$Z_1 = P_1^\perp a_1 \otimes a_2 P_2, \quad Z_2 = P_1^\perp \otimes a_2 P_2^\perp, \quad Z_3 = P_1 a_1 \otimes P_2, \quad Z_4 = P_1 \otimes P_2^\perp.$$

In order to compute the ‘first-return moments’ $N_Z(n)$ in the state φ , we need to compute the mixed φ -moments of Z_i ’s. However, it turns out that there are only two types of them which give a non-vanishing contribution. We compute them in Propositions 4.3–4.4.

Proposition 4.2. *For r odd and $j_1, j_2, \dots, j_r \geq 1$, we have*

$$\varphi(Z_3 Z_1^{j_1} Z_2^{j_2} Z_1^{j_3} \dots Z_1^{j_r}) = N_{a_1}(m) N_{a_2}(k_1) \dots N_{a_2}(k_{m-1}),$$

where $m = 1 + \sum_{k \text{ odd}} j_k$ and all k_i 's are equal to 1, except

$$k_{j_1+1} = j_2 + 1, \quad k_{j_1+j_3+1} = j_4 + 1, \dots, k_{j_1+j_3+\dots+j_{r-2}+1} = j_{r-1} + 1,$$

and where the 'first return moments' $N_{a_\iota}(s)$ refer to the state φ_ι , $\iota \in I$.

Proof. We have

$$\begin{aligned} \varphi(Z_3 Z_1^{j_1} Z_2^{j_2} Z_1^{j_3} \dots Z_1^{j_r}) &= \varphi_1(P_1 a_1 (P_1^\perp a_1)^{j_1} P_1^\perp (P_1^\perp a_1)^{j_3} \dots P_1^\perp (P_1^\perp a_1)^{j_r}) \\ &\quad \varphi_2(P_2 (a_2 P_2)^{j_1} (a_2 P_2^\perp)^{j_2} (a_2 P_2)^{j_3-1} \dots (a_2 P_2)^{j_r}) \\ &= \varphi_1(a_1 (P_1^\perp a_1)^{j_1+j_3+\dots+j_r}) \\ &\quad \varphi_2((a_2 P_2)^{j_1} (a_2 P_2^\perp)^{j_2} (a_2 P_2)^{j_3} \dots (a_2 P_2)^{j_r}) \\ &= N_{a_1}(1 + j_1 + j_3 + \dots + j_r) (N_{a_2}(1))^{j_1+(j_3-1)+\dots+(j_r-1)} \\ &\quad N_{a_2}(j_2 + 1) N_{a_2}(j_4 + 1) \dots N_{a_2}(j_{r-1} + 1), \end{aligned}$$

using the following properties:

$$P_1 a_1 P_1 = \varphi_1(a_1) P_1 = N_{a_1}(1) P_1, \quad P_2 a_2 P_2 = \varphi_2(a_2) P_2 = N_{a_2}(1) P_2.$$

Setting $k_1 = k_2 = \dots = k_{j_1} = 1$, $k_{j_1+1} = j_2 + 1$, $k_{j_1+2} = \dots = k_{j_1+j_3} = 1$, $k_{j_1+j_3} = 1$, $k_{j_1+j_3+1} = j_4 + 1$, $\dots$, $k_{j_1+j_3+\dots+j_{r-2}+1} = j_{r-1} + 1$, $k_{j_1+j_3+\dots+j_{r-2}+2} = \dots = k_{m-1} = 1$, we get the desired result. $\blacksquare$

Proposition 4.3. For r even and $j_1, j_2, \dots, j_r \geq 1$, we have

$$\varphi(Z_3 Z_2^{j_1} Z_1^{j_2} Z_2^{j_3} \dots Z_1^{j_r}) = N_{a_1}(m) N_{a_2}(k_1) \dots N_{a_2}(k_{m-1}),$$

where $m = 1 + \sum_{k \text{ even}} j_k$ and all k_i 's are equal to 1, except

$$k_{j_2+1} = j_3 + 1, \quad k_{j_2+j_4+1} = j_5 + 1, \dots, k_{j_2+j_4+\dots+j_{r-2}+1} = j_{r-1} + 1,$$

and where the 'first return moments' $N_{a_\iota}(s)$ refer to the state φ_ι , $\iota \in I$.

Proof. Using similar arguments as in the proof of Proposition 4.3, we get

$$\begin{aligned} \varphi(Z_3 Z_2^{j_1} Z_1^{j_2} Z_2^{j_3} \dots Z_1^{j_r}) &= \varphi_1(P_1 a_1 P_1^\perp (P_1^\perp a_1)^{j_2} P_1^\perp (P_1^\perp a_1)^{j_4} \dots (P_1^\perp a_1)^{j_r}) \\ &\quad \varphi_2(P_2 (a_2 P_2^\perp)^{j_1} (a_2 P_2)^{j_2} (a_2 P_2^\perp)^{j_3} \dots (a_2 P_2)^{j_r}) \\ &= \varphi_1(a_1 (P_1^\perp a_1)^{j_2+j_4+\dots+j_r}) \\ &\quad \varphi_2((a_2 P_2^\perp)^{j_1} (a_2 P_2)^{j_2} (a_2 P_2^\perp)^{j_3} \dots (a_2 P_2)^{j_r}) \\ &= N_{a_1}(1 + j_2 + j_4 + \dots + j_r) (N_{a_2}(1))^{(j_2-1)+(j_4-1)+\dots+(j_r-1)} \\ &\quad N_{a_2}(j_1 + 1) N_{a_2}(j_3 + 1) \dots N_{a_2}(j_{r-1} + 1). \end{aligned}$$

Setting $k_1 = j_1 + 1$, $k_2 = \dots = k_{j_2} = 1$, $k_{j_2+1} = j_3 + 1$, $k_{j_2+2} = \dots = k_{j_2+j_4} = 1$, $k_{j_2+j_4+1} = j_5 + 1$, $\dots$, $k_{j_2+j_4+\dots+j_{r-2}+1} = j_{r-1} + 1$, $k_{j_2+j_4+\dots+j_{r-2}+2} = \dots = k_{m-1} = 1$, we get the desired result. $\blacksquare$

Theorem 4.1. If the φ -distribution of a_2 is not concentrated at zero, then the formal power series corresponding to random variables a_1, a_2 and Z satisfy the relation

$$(4.4) \quad \eta_Z(z) = \frac{z \eta_{a_1}(\eta_{a_2}(z))}{\eta_{a_2}(z)}.$$

If the distribution of a_2 is concentrated at zero, then $\eta_Z(z) = z\mu_1(X)$, where μ_1 is the φ -distribution of a_1 .

Proof. Suppose the φ -distribution of a_2 is not the Dirac delta. Then, writing the RHS of equation (4.4) as $D(z) = \sum_{n=1}^{\infty} d_n z^n$, we need to show that $d_n = \eta_Z(n)$ for $n \in \mathbb{N}$. We clearly have

$$N_Z(1) = \varphi(Z_3) = \varphi_1(a_1) = N_{a_1}(1) = d_1.$$

Now, for $n > 1$, it holds that

$$d_n = \sum_{m=1}^n N_{a_1}(m) \sum_{k_1+k_2+\dots+k_{m-1}=n-1} N_{a_1}(k_1)N_{a_2}(k_2)\dots N_{a_2}(k_{m-1}),$$

where all k_i 's are assumed to be positive integers, Using Proposition 2.2 and the explicit form of Z_i 's, we get

$$\begin{aligned} N_Z(n) &= \varphi(Z(P^\perp Z)^{n-1}) \\ &= \varphi(Z_3(Z_1 + Z_2)^{n-1}) \\ &= \varphi(Z_3(Z_1 + Z_2)^{n-2}Z_1) \\ &= \sum_{r=1}^{n-2} \sum_{i_1 \neq i_2 \neq \dots \neq i_r} \sum_{j_1+j_2+\dots+j_r=n-2} \varphi(Z_3 Z_{i_1}^{j_1} Z_{i_2}^{j_2} \dots Z_{i_r}^{j_r} Z_1), \end{aligned}$$

where $j_1, j_2, \dots, j_r$ are positive integers. Note that on the RHS of the formula for d_n we have the sum of products

$$N_{a_1}(m)N_{a_2}(k_1)N_{a_2}(k_2)\dots N_{a_2}(k_{m-1}),$$

where $1 \leq m \leq n$ and $(k_1, k_2, \dots, k_{m-1})$ is a sequence of positive integers which add up to $n - 1$. Now, every product obtained in Propositions 4.3-4.4 is of exactly this form, with indices $(k_1, k_2, \dots, k_{m-1})$ satisfying the conditions just mentioned. Moreover, all these products are different from each other and exhaust all possible values of $(m, k_1, \dots, k_{m-1})$. In fact, there are two disjoint cases distinguished: Proposition 4.3 covers the case $k_1 = 1$ and Proposition 4.4 – the case $k_1 \neq 1$. In the first case, even j_i 's determine the subsequence of $(k_1, k_2, \dots, k_{m-1})$ consisting of numbers which are greater than 1 and odd j_i 's determine their positions in the sequence by determining all k_i 's which are equal to 1. In the second case, odd j_i 's determine the subsequence of $(k_1, k_2, \dots, k_{m-1})$ consisting of numbers which are greater than 1 and even j_i 's determine their positions in the sequence by determining all k_i 's which are equal to 1. Therefore, both cases give together every tuple $(m, k_1, k_2, \dots, k_{m-1})$ under consideration exactly once. This proves (4.4) in the case when the distribution of a_2 is not concentrated at zero. Finally, if the φ -distribution of a_2 is δ_0 , then $\varphi_2(a_2^n) = \delta_{n,0}$, which easily gives $N_Z(n) = N_{a_1}(1)\delta_{n,1}$, where $\delta_{i,j}$ is the Kronecker symbol, and since $N_{a_1}(1) = \varphi_1(a_1) = \mu_1(X)$, we have $\eta_Z(z) = z\mu_1(X)$. This completes the proof. $\blacksquare$

This shows that Eq.(1.7) holds for the orthogonal multiplicative convolution of distributions $\mu_1, \mu_2 \in \Sigma$, provided $\mu_2 \neq \delta_0$, where the functions η are treated as formal

power series. In turn, explicit computations of distributions $\mu_1 \angle \mu_2$ based on Definition 4.2 give

$$(4.5) \quad \delta_0 \angle \mu = \delta_0 \quad \text{and} \quad \mu \angle \delta_0 = \delta_{\mu(X)}$$

for any $\mu \in \Sigma$. Let us also remark that Theorem 4.5 gives another formula for formal power series corresponding to φ -distributions, namely

$$(4.6) \quad \rho_{\mu_1 \angle \mu_2}(z) = \rho_{\mu_1}(\eta_{\mu_2}(z)),$$

where ρ_{μ_1} is given by (3.3), which turns out slightly more useful in computations involving $\mu_1 \angle \mu_2$ than (4.4) (see Section 5).

Corollary 4.2. *If μ_1 and μ_2 are φ -distributions of certain random variables, then the φ -moment of $\mu_1 \angle \mu_2$ of order $n \in \mathbb{N}$ depends on φ -moments of μ_1 of orders $1 \leq k \leq n$ and φ -moments of μ_2 of orders $1 \leq k \leq n - 1$.*

Proof. This fact is a consequence of Proposition 4.1 and Theorem 4.5. In fact, the combinatorial formula for d_n in the proof of Theorem 4.5 gives the explicit formula for $N_{\mu \angle \nu}(n)$ in terms of $N_\mu(m)$ and $N_\nu(k)$ for $1 \leq m \leq n$ and $1 \leq k \leq n - 1$. Thus, in view of (1.2), a similar property holds for the φ -moments of $\mu \angle \nu$. ■

Example 4.1. If $\mu_1, \mu_2 \in \Sigma$, the lowest order first return moments $N_{\mu_1 \angle \mu_2}(n)$ are given by

$$\begin{aligned} N_{\mu_1 \angle \mu_2}(1) &= N_{\mu_1}(1), \\ N_{\mu_1 \angle \mu_2}(2) &= N_{\mu_1}(2)N_{\mu_2}(1), \\ N_{\mu_1 \angle \mu_2}(3) &= N_{\mu_1}(3)N_{\mu_2}^2(1) + N_{\mu_1}(2)N_{\mu_2}(2), \\ N_{\mu_1 \angle \mu_2}(4) &= N_{\mu_1}(4)N_{\mu_2}^3(1) + 2N_{\mu_1}(3)N_{\mu_2}(2)N_{\mu_2}(1) + N_{\mu_1}(2)N_{\mu_2}(3). \end{aligned}$$

When we return to the φ -moments $\mu_1 \angle \mu_2(X^n)$, using (1.2), we can express them in terms of ‘universal polynomials’ $Q_n(\mu_1(X), \dots, \mu_1(X^n), \mu_2(X), \dots, \mu_2(X^{n-1}))$, but one should note that they are not homogenous as in the case of free convolutions [25].

5. CONVOLUTIONS OF MEASURES ON $\mathbb{R}_+$ AND $\mathbb{T}$.

In this section we show that the orthogonal multiplicative convolution can be defined for arbitrary probability measures on $\mathbb{R}_+$ which are not concentrated at zero, as well as for arbitrary probability measures on the unit circle $\mathbb{T}$.

For the following result, we refer the reader to the works of Belinschi and Bercovici [3,4].

Theorem 5.1. *There is a bijection between $\mathcal{M}_{\mathbb{R}_+}$ and the class of analytic self-maps η of $\mathbb{C} \setminus \mathbb{R}_+$, such that*

- (1) $\eta(\bar{z}) = \overline{\eta(z)}$ for all $z \in \mathbb{C} \setminus \mathbb{R}_+$.
- (2) $\lim_{z \rightarrow 0^-, z < 0} \eta(z) = 0$
- (3) $\arg \eta(z) \in [\arg z, \pi)$ for all $z \in \mathbb{C}^+$.

Moreover, the map η corresponding to $\mu \in \mathcal{M}_{\mathbb{R}_+}$ is given by the transform η_μ .

We shall use this result to prove that formula (4.4), which was shown to hold for formal power series corresponding to distributions of random variables, can also be used to define $\mu_1 \angle \mu_2$ for $\mu_1, \mu_2 \in \mathcal{M}_{\mathbb{R}_+}$ and $\mu_2 \neq \delta_0$, if formal power series are replaced by analytic functions on $\mathbb{C} \setminus \mathbb{R}_+$.

Proposition 5.1. *If $\mu_1, \mu_2 \in \mathcal{M}_{\mathbb{R}_+}$ and μ_2 is not concentrated at zero, then there exists a unique probability measure $\mu \in \mathcal{M}_{\mathbb{R}_+}$, such that*

$$(5.1) \quad \eta_\mu(z) = \frac{z\eta_{\mu_1}(\eta_{\mu_2}(z))}{\eta_{\mu_2}(z)}$$

for $z \in \mathbb{C} \setminus \mathbb{R}_+$. The measure μ will be defined to be the orthogonal multiplicative convolution of μ_1 and μ_2 , denoted $\mu_1 \angle \mu_2$.

Proof. For $z \in \mathbb{C} \setminus \mathbb{R}_+$, let us define

$$\eta(z) = \frac{z\eta_{\mu_1}(\eta_{\mu_2}(z))}{\eta_{\mu_2}(z)}.$$

Since $\mu_1, \mu_2 \in \mathcal{M}_{\mathbb{R}_+}$, transforms η_{μ_1} and η_{μ_2} are analytic self-maps of $\mathbb{C} \setminus \mathbb{R}_+$ which satisfy the conditions of Theorem 5.1. This implies that $\eta(z)$ is an analytic self-map of $\mathbb{C} \setminus \mathbb{R}_+$ since $\mu_2 \neq \delta_0$. Condition (1) of Theorem 5.1 holds for η_{μ_1} and η_{μ_2} and therefore it holds for η . Then, it is known that

$$0 \leq \lim_{z \rightarrow 0^-, z < 0} \frac{\eta_{\mu_2}(z)}{z} \leq \infty,$$

and this limit is different from zero (it equals zero only if $\mu_2 = \delta_0$). Thus, we obtain

$$\begin{aligned} \lim_{z \rightarrow 0^-, z < 0} \eta(z) &= \lim_{z \rightarrow 0^-, z < 0} \frac{z}{\eta_{\mu_2}(z)} \lim_{z \rightarrow 0^-, z < 0} \eta_{\mu_1}(\eta_{\mu_2}(z)) \\ &= c \cdot \lim_{z \rightarrow 0^-, z < 0} \eta_{\mu_1}(\eta_{\mu_2}(z)) \\ &= 0 \end{aligned}$$

for some $0 \leq c < \infty$, where we used condition 2 of Theorem 5.1 for the function η_{μ_1} (if $z \rightarrow 0^-$, $\psi_{\mu_2}(z) \rightarrow 0^-$ and thus $\eta_{\mu_2}(z) \rightarrow 0^-$ and therefore $\eta_{\mu_1}(\eta_{\mu_2}(z)) \rightarrow 0$ as $z \rightarrow 0^-$). This proves that condition (2) of Theorem 5.1 holds for the map η .

Next, for any $z \in \mathbb{C}^+$, we have $\psi_{\mu_2}(z) \in \mathbb{C}^+$, using the integral representation of $\psi_{\mu_2}(z)$ given by (1.1). This implies that $\eta_{\mu_2}(z) \in \mathbb{C}^+$, since

$$\eta_{\mu_2}(z) = \frac{\psi_{\mu_2}(z)}{1 + \psi_{\mu_2}(z)} = \frac{\psi_{\mu_2}(z) + |\psi_{\mu_2}(z)|^2}{1 + |\psi_{\mu_2}(z)|^2},$$

and therefore

$$\Im \eta_{\mu_2}(z) = \frac{\Im \psi_{\mu_2}(z)}{|1 + \psi_{\mu_2}(z)|^2}.$$

Hence, applying Theorem 5.1 to the function η_{μ_1} , we get

$$\arg \eta_{\mu_1}(\eta_{\mu_2}(z)) \geq \arg \eta_{\mu_2}(z),$$

and thus

$$\arg \eta(z) = \arg z + \arg \eta_{\mu_1}(\eta_{\mu_2}(z)) - \arg \eta_{\mu_2}(z) \geq \arg z.$$

Similarly,

$$\begin{aligned} \arg \eta(z) &= \arg \eta_{\mu_1}(\eta_{\mu_2}(z)) + \arg z - \arg \eta_{\mu_2}(z) \\ &= \arg \eta_{\mu_1}(\eta_{\mu_2}(z)) - (\arg \eta_{\mu_2}(z) - \arg z) \\ &< \pi \end{aligned}$$

since we have

$$\arg \eta_{\mu_2}(z) \geq \arg z \quad \text{and} \quad \arg \eta_{\mu_1}(\eta_{\mu_2}(z)) < \pi,$$

which follows from $\eta_{\mu_2}(z) \in \mathbb{C}^+$. This proves that η satisfies condition (3) of Theorem 5.1, which completes the proof. $\blacksquare$

For computations, it is convenient to use (4.6) and represent the transforms involved as continued fractions. The continued fraction representation of η_μ for compactly supported $\mu \in \mathcal{M}_{\mathbb{R}_+}$ is obtained from that of the Cauchy transform G_μ , or its reciprocal F_μ , by using the formula

$$(5.2) \quad \eta_\mu(z) = 1 - zF_\mu\left(\frac{1}{z}\right).$$

If $\mu \in \mathcal{M}_{\mathbb{R}_+}$ is associated with sequences of Jacobi parameters (α, ω) , where $\alpha = (\alpha_n)_{n \geq 0}$ and $\omega = (\omega_n)_{n \geq 0}$, which we write $J(\mu) = (\alpha, \omega)$, then, using the continued fraction representation of G_μ , we obtain

$$(5.3) \quad \eta_\mu(z) = \alpha_0 z + \frac{\omega_0 z^2}{1 - \alpha_1 z - \frac{\omega_1 z^2}{1 - \alpha_2 z - \frac{\omega_2 z^2}{1 - \alpha_3 z - \frac{\omega_3 z^2}{\dots}}}}$$

and a related continued fraction for ρ_μ . Conversely, this formula, together with the first formula of (1.7), enables us to compute the Jacobi sequences corresponding to the orthogonal multiplicative convolution. In the case of compactly supported measures, the Jacobi sequences uniquely determine the corresponding measure.

Example 5.1. Let δ_a be the Dirac measure with $a > 0$ and let $\mu \in \mathcal{M}_{\mathbb{R}_+}$ be compactly supported, with the associated Jacobi sequences $J(\mu) = (\alpha, \omega)$. Then $\rho_\mu(z) = a$ and therefore, $\rho_{\delta_a \angle \mu}(z) = \rho_{\delta_a}(\eta_\mu(z)) = a$, which gives $\delta_a \angle \mu = \delta_a$. In turn, $\rho_{\mu \angle \delta_a}(z) = \rho_\mu(az)$. Let us denote the corresponding transformation of compactly supported measures by S_a . In terms of reciprocal Cauchy transforms, it can be defined by a convex linear combination

$$F_{S_a \mu}(z) = \frac{1}{a} F_{D_a \mu}(z) + \left(1 - \frac{1}{a}\right) z,$$

where $D_\lambda \mu$ is the dilation of measure μ by λ , defined by $D_\lambda \mu(E) = \mu(\lambda^{-1}E)$. In terms of the transforms η , we have $\eta_{S_a \mu}(z) = \eta_{D_a \mu}(z)/a$, using (5.2). We also have $\eta_{D_a \mu}(z) = \eta_\mu(az)$. Note that $S_a \mu = (T_a \circ D_a) \mu$, where $T_{t^{-1}}$ is the so-called t -transformation of measures [9]. If μ is compactly supported, we can write $\eta_{S_a \mu}$ in the form of a continued

fraction

$$\eta_{S_a\mu}(z) = \alpha_0 z + \frac{\omega_0 a z^2}{1 - \alpha_1 a z - \frac{\omega_1 a^2 z^2}{1 - \alpha_2 a z - \frac{\omega_2 a^2 z^2}{1 - \alpha_3 a z - \frac{\omega_3 a^2 z^2}{\ddots}}}}.$$

In particular, $\mu \angle \delta_1 = \mu$, thus δ_1 is the right unit w.r.t. the operation $\angle$ on $\mathcal{M}_{\mathbb{R}_+}$, and $\delta_1 \angle \mu = \delta_1$.

Example 5.2. If $\mu_1 = (1-p)\delta_0 + p\delta_1$ and $\mu_2 = (1-q)\delta_0 + q\delta_1$, then the corresponding reciprocal Cauchy transforms are

$$F_{\mu_1}(z) = \frac{z-1+p}{z(z-1)}, \quad F_{\mu_2}(z) = \frac{z-1+q}{z(z-1)},$$

which gives

$$\rho_{\mu_1} = p + \frac{(p-p^2)z}{1-(1-p)z}, \quad \eta_{\mu_2} = qz + \frac{(q-q^2)z^2}{1-(1-q)z},$$

and therefore, in view of (4.6), we obtain

$$\rho_{\mu_1 \angle \mu_2}(z) = p + \frac{(p-p^2)qz}{1+(pq-1)z}.$$

Using (5.3), we obtain $J(\mu_1 \angle \mu_2) = (\alpha, \omega)$, with $\alpha = (p, 1-pq, 0, 0, \dots)$ and $\omega = ((p-p^2)q, 0, 0, \dots)$. The corresponding measure has two atoms (explicit dependence on p and q is rather complicated and is omitted).

A result analogous to Theorem 5.1 also holds for the set $\mathcal{M}_{\mathbb{T}}$ of probability measures on the unit circle $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$. Here, the class of self-maps η of the open unit disc $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ is used, which is again related to the transforms η_μ . For details, we refer the reader to the works of Belinschi and Bercovici [3,4].

Theorem 5.2. *There is a bijection between $\mathcal{M}_{\mathbb{T}}$ and the class of analytic self-maps η of $\mathbb{D}$ such that $\eta(0) = 0$ and $|\eta(z)| \leq |z|$ for all $z \in \mathbb{D}$. Moreover, the map η corresponding to μ is given by the transform η_μ .*

Proposition 5.2. *If $\mu_1, \mu_2 \in \mathcal{M}_{\mathbb{T}}$, then there exists a unique probability measure $\mu \in \mathcal{M}_{\mathbb{T}}$ such that*

$$(5.4) \quad \eta_\mu(z) = \frac{z\eta_{\mu_1}(\eta_{\mu_2}(z))}{\eta_{\mu_2}(z)}$$

for $z \in \mathbb{D}$, where we understand that $\eta_{\mu_1}(0)/0 = \eta'_{\mu_1}(0)$. This measure μ will be defined to be the orthogonal multiplicative convolution $\mu_1 \angle \mu_2$.

Proof. First of all, observe that $|\eta(z)| \leq |z|$ since η_{μ_1} has the same property by Theorem 5.3. Moreover, η is a quotient of analytic functions on $\mathbb{D}$, and if $\eta_{\mu_2}(z_0) = 0$, then

$$\eta(z_0) = \lim_{z \rightarrow z_0} z \frac{\eta_{\mu_1}(\eta_{\mu_2}(z))}{\eta_{\mu_2}(z)} = z_0 \eta'_{\mu_1}(0),$$




FIGURE 3. An example of the orthogonal loop product

thus η is a well-defined analytic self-map of $\mathbb{D}$. Finally, $\eta(0) = 0 \cdot \eta'_{\mu_1}(0) = 0$. Thus, η satisfies the conditions of Theorem 5.3 and therefore $\eta = \eta_\mu$ for some $\mu \in \mathcal{M}_{\mathbb{T}}$, which completes the proof. $\blacksquare$

Example 5.3. For any $\mu \in \mathcal{M}_{\mathbb{T}}$ and $a \in \mathbb{T}$, we get $\delta_a \angle \mu = \delta_a$ and $\mu \angle \delta_a = S_a \mu$, where $S_a \mu$ is defined by the same equation $\eta_{S_a \mu}(z) = \eta_{D_a \mu}(z)/a$ as in the case of measures on $\mathbb{R}_+$ (on $\mathcal{M}_{\mathbb{T}}$, the operation D_a should be interpreted as a rotation).

6. ORTHOGONAL LOOP PRODUCT OF GRAPHS

In this section we establish a relation between the orthogonal multiplicative convolution of compactly supported probability measures and a new type of product of graphs called the ‘orthogonal loop product’.

As in the monotone case, we begin with recalling the definition of the orthogonal product of graphs [2,16]. Then we will modify this product in an appropriate manner to find a relation with the orthogonal multiplicative convolution.

Definition 6.1. The *orthogonal product* of two rooted graphs $(\mathcal{G}_1, e_1)$ and $(\mathcal{G}_2, e_2)$ is the rooted graph $(\mathcal{G}_1 \vdash \mathcal{G}_2, e)$ obtained by attaching a copy of $\mathcal{G}_2$ by its root e_2 to each vertex of $\mathcal{G}_1$ but the root e_1 , where e is taken to be equal to e_1 . If its set of vertices is identified with $V_1 \vdash V_2 := (V_1^0 \times V_2) \cup \{e_1 \times e_2\}$, then e is identified with $e_1 \times e_2$.

It is worth noting that the orthogonal product of graphs resembles their comb product. The difference is that in the comb product the second graph is glued by its root to *all* vertices of the first graph, whereas in the orthogonal product the second graph is glued to *all vertices but the root* of the first graph.

Definition 6.2. Suppose the edges of the orthogonal product $(\mathcal{G}_1 \vdash \mathcal{G}_2, e)$ are I -colored. The *orthogonal loop product* of rooted graphs $(\mathcal{G}_1, e_1)$ and $(\mathcal{G}_2, e_2)$ is the rooted graph $(\mathcal{G}_1 \vdash_\ell \mathcal{G}_2, e)$ obtained from $(\mathcal{G}_1 \vdash \mathcal{G}_2, e)$ by attaching a loop of color 1 to all vertices but the root of each copy of $\mathcal{G}_2$, and a loop of color 2 to the root of $\mathcal{G}_1$.

The justification of the above definition is similar to that in Remark 3.1. It is enough to deduce the glueing rules from the expressions for R_1 and R_2 , given in the proof of Proposition 4.1, interpreting a_ι as the adjacency matrix of graph $\mathcal{G}_\iota$, $\iota \in I$. Also, there is an easy analog of Remark 3.1. Finally, it follows from [2] that the additive analog of the Multiplication Theorem holds for orthogonal independence. An example of the orthogonal loop product of graphs is given in Fig.3 (cf. Figs.1-2).

Example 6.1. In the orthogonal loop product of graphs given in Fig.3, we have

$$(\beta_1, w_1, \beta_2, w_2, \beta_3, w_3, \beta_4, w_4) \in D_8(e),$$

where the sequence of edges of color 1 is the same as in Example 3.1, namely

$$(\beta_1, \beta_2, \beta_3, \beta_4) = ((e, x), (x, x'), (x', x), (x, e)),$$

and the (alternating) f-walks attached to the vertices x, x', x, e (in that order) are: $w_1 = w_3 = (x, x)_2$, $w_2 = (x', x')_2$ and $w_4 = (e, e)$. Note that this d-walk has length 8. Again, as in the case of d-walks on the comb loop product, if any of the loops at x or x' is replaced by an alternating f-walk, we shall still get a rooted alternating d-walk. Of course, the simplest rooted alternating d-walk is given by $(\beta_1, w_1, \beta_4, w_4) \in D_4(e)$ and $(\beta_0, w_4) \in D_2(e)$, where $\beta_0 = (e, e)_1$.

Theorem 6.1. *The Multiplication Theorem holds for orthogonal independence, the associated loop product $\mathcal{G}_1 \vdash_\ell \mathcal{G}_2$, and the multiplicative convolution $\mu_1 \angle \mu_2$.*

Proof. The proof is similar to that of Theorem 3.1, but the combinatorics is based on a different formula for formal power series, namely

$$\eta_{\mu_1 \angle \mu_2}(z) = \frac{z \eta_{\mu_1}(\eta_{\mu_2}(z))}{\eta_{\mu_2}(z)},$$

which leads to the combinatorial formula

$$N_{\mu_1 \angle \mu_2}(n) = \sum_{r=1}^n N_{\mu_1}(r) \sum_{k_1+k_2+\dots+k_{r-1}=n-1} N_{\mu_2}(k_1) N_{\mu_2}(k_2) \dots N_{\mu_2}(k_{r-1}),$$

where it is assumed that $k_1, k_2, \dots, k_{r-1}$ are positive integers. Recall that in the orthogonal product $\mathcal{G}_1 \vdash \mathcal{G}_2$ there is one copy of $\mathcal{G}_1$ (with e_1 identified with the root e of the product graph) with a copy of $\mathcal{G}_2$ attached by its root to every vertex x of $\mathcal{G}_1$ but the root e_1 . Therefore, each d-walk $w \in D_{2n}(e)$ is a sequence of r edges of $\mathcal{G}_1$, which themselves must form an f-walk $c = (e, v_1, v_2, \dots, v_{r-1}, e)$ of color 1, interlaced with alternating f-walks $w_i \in F(v_i)$, $1 \leq i \leq r-1$ and a loop of color 2, $w_r = (e, e)_2$. Note that this is the only way to produce a rooted alternating d-walk since the only edge of color 2 incident on e is the loop and therefore, in order to get an alternating walk, the first f-walk which begins and ends with an edge of color 1 must be followed by the loop at e of color 2 to make a double return to e . The contribution from each product of type

$$N_{\mu_1}(r) N_{\mu_2}(k_1) N_{\mu_2}(k_2) \dots N_{\mu_2}(k_{r-1})$$

to the RHS of the above formula is equal to the number of all such f-walks c of color 1 which have r edges and are interlaced with $r-1$ f-walks of lengths $2k_1-1, 2k_2-1, \dots, 2k_{r-1}-1$ attached to all vertices of c but one (we choose this vertex to be the root since we want the considered walk to be an f-walk). The summation over all $1 \leq r \leq n$ and $2k_1 + 2k_2 + \dots + 2k_{r-1} - r + 1 = 2n - 1$ indicates that the first f-walk u_1 in the d-walk $w = (u_1, u_2)$ is of length $2n-1$, and u_2 is the loop of length 1. The summation over $1 \leq r \leq n$ gives exactly the cardinality of $D_{2n}(e)$, which finishes the proof. $\blacksquare$

Example 6.2. Let us apply the formulas of Example 4.1 to the enumeration of rooted alternating d-walks on the graph $\mathcal{G}_1 \vdash_\ell \mathcal{G}_2$ in Fig.3 (we keep the notation of Example 3.2 for spectral distributions μ_1 and μ_2 and thus we get the same values of the $N_{\mu_1}(k)$'s and the $N_{\mu_2}(j)$'s). Using Example 4.1 and Theorem 6.1, we get $D_2(e) = D_4(e) = D_6(e) = D_8(e) = 1$, which can be verified directly by counting rooted alternating d-walks on $\mathcal{G}_1 \vdash_\ell \mathcal{G}_2$.

7. SUBORDINATION OPERATORS

In analogy to the additive case [16], where we introduced and studied operators related to the subordination property for the free additive convolution, we shall now present an analogous approach to the subordination property for the free multiplicative convolution. We will mainly refer to sets $\mathcal{M}_{\mathbb{R}^+}$ and $\mathcal{M}_{\mathbb{T}}$, which correspond to positive and unitary operators, but the operatorial subordination will also hold for all bounded operators.

Let $(\mathcal{H}_\iota, \xi_\iota)$, where $\iota \in I$, be Hilbert spaces with distinguished unit vectors. Then their Hilbert space free product $(\mathcal{H}_1, \xi_1) * (\mathcal{H}_2, \xi_2)$ is $(\mathcal{H}, \xi)$ where

$$(7.1) \quad \mathcal{H} = \mathbb{C} \xi \oplus \bigoplus_{n=1}^{\infty} \bigoplus_{\iota_1 \neq \iota_2 \neq \dots \neq \iota_n} \mathcal{H}_{\iota_1}^0 \otimes \mathcal{H}_{\iota_2}^0 \otimes \dots \otimes \mathcal{H}_{\iota_n}^0,$$

with $\mathcal{H}_\iota^0 = \mathcal{H}_\iota \ominus \mathbb{C} \xi_\iota$ and ξ denoting a unit vector (canonical scalar product is used). For any $h \in \mathcal{H}_\iota$, denote by h^0 the orthogonal projection of h onto $\mathcal{H}_\iota^0$. Moreover, let

$$(7.2) \quad \mathcal{H}^{(n)}(\iota) = \bigoplus_{\substack{\iota_1 \neq \iota_2 \neq \dots \neq \iota_n \\ \iota_1 \neq j}} \mathcal{H}_{\iota_1}^0 \otimes \mathcal{H}_{\iota_2}^0 \otimes \dots \otimes \mathcal{H}_{\iota_n}^0,$$

$$(7.3) \quad \mathcal{K}^{(n)}(\iota) = \bigoplus_{\substack{\iota_1 \neq \iota_2 \neq \dots \neq \iota_n \\ \iota_n \neq \iota}} \mathcal{H}_{\iota_1}^0 \otimes \mathcal{H}_{\iota_2}^0 \otimes \dots \otimes \mathcal{H}_{\iota_n}^0,$$

for $\iota \in I$ and $n \in \mathbb{N}$, and, for convenience, we set $\mathcal{H}^{(0)}(\iota) = \mathcal{K}^{(0)}(\iota) = \mathbb{C} \xi$ with the canonical projection $P_\xi : \mathcal{H} \rightarrow \mathbb{C} \xi$.

Since our index set I consists of two elements, the above notation gives identifications

$$(7.4) \quad \mathcal{H}^{(n)}(\iota) = \begin{cases} \mathcal{K}^{(n)}(\iota) & n \text{ odd,} \\ \mathcal{K}^{(n)}(\bar{\iota}) & n \text{ even.} \end{cases}$$

Nevertheless, it is convenient to use both sequences, (7.2) and (7.3), as well as direct sums

$$(7.5) \quad \mathcal{H}(\iota) = \bigoplus_{n=1}^{\infty} \mathcal{H}^{(n)}(\iota) \quad \text{and} \quad \mathcal{K}(\iota) = \bigoplus_{n=1}^{\infty} \mathcal{K}^{(n)}(\iota),$$

where $\iota \in I$.

As in our previous work, we decompose the free product of Hilbert spaces as (two different) orthogonal direct sums

$$(7.6) \quad \mathcal{H} = \bigoplus_{n=1}^{\infty} \mathcal{H}^{(n-1)}(\iota) \oplus \mathcal{H}^{(n)}(\bar{\iota}),$$

where $\iota \in I$, and denote by

$$(7.7) \quad P_\iota(n) : \mathcal{H} \rightarrow \mathcal{H}^{(n-1)}(\iota) \oplus \mathcal{H}^{(n)}(\bar{\iota})$$

the associated canonical projections. Finally, we define the so-called *vacuum state* $\varphi(\cdot) = \langle \cdot, \xi, \xi \rangle$ on $\mathcal{B}(\mathcal{H})$.

Of particular interest will be *s-free products of Hilbert spaces*, $(\mathcal{H}_\iota, \xi_\iota)$ and $(\mathcal{H}_{\bar{\iota}}, \xi_{\bar{\iota}})$, denoted $(\mathcal{H}_\iota \oplus \mathcal{H}_{\bar{\iota}}, \xi)$ and defined [16] as the pair $(\mathcal{K}_\iota, \xi)$ where

$$(7.8) \quad \mathcal{K}_\iota = \mathbb{C}\xi \oplus \mathcal{K}(\bar{\iota}),$$

and $\iota \in I$. The direct sum decompositions

$$(7.9) \quad \mathcal{K}_\iota = \bigoplus_{n \text{ odd}} \mathcal{H}^{(n-1)}(\iota) \oplus \mathcal{H}^{(n)}(\bar{\iota})$$

hold for each $\iota \in I$.

Let $x \in B(\mathcal{H}_1)$ and $y \in B(\mathcal{H}_2)$ be fixed random variables. The corresponding free random variables $X_1 := \lambda(x)$ and $X_2 := \lambda(y)$ are elements of $B(\mathcal{H})$, where λ is the free product representation on $\mathcal{H}$, and can be decomposed according to (7.6) as

$$(7.10) \quad X_\iota = \sum_{j=1}^{\infty} X_\iota(j),$$

where $X_\iota(j) = P_\iota(j)X_\iota P_\iota(j)$, $j \in \mathbb{N}$, can be viewed as replicas of x and y , respectively, where $\iota \in I$. Using (7.6) and (7.9), we also have the decompositions

$$(7.11) \quad 1 = \sum_{j=1}^{\infty} P_\iota(j) \quad \text{and} \quad 1_{\mathcal{K}_\iota} = \sum_{j \text{ odd}} P_\iota(j)$$

of the units in $B(\mathcal{H})$ and $B(\mathcal{K}_\iota)$, where $\iota \in I$, respectively. Here, and also in the sequel, we denote by the same symbols, $X_\iota(j)$ and $P_\iota(j)$, the corresponding operators on $\mathcal{K}_\iota$.

We shall use representations of free random variables as ‘orthogonal series’ with the unit singled out, namely

$$(7.12) \quad X_\iota = 1 + \sum_{j=1}^{\infty} x_\iota(j)$$

where $x_\iota(j) = X_\iota(j) - P_\iota(j)$ and $\iota \in I$, $j \in \mathbb{N}$. If no confusion arises, we will distinguish the first term in the above series by a special notation and write $x_\iota = x_\iota(1)$. Let us remark that the above form is suitable for the study of multiplicative convolutions, where units play a special role (roughly speaking, they have to be subtracted when it comes to proving some kind of independence).

We begin with the decomposition of the product $X_{\bar{\iota}}X_\iota$ which corresponds to Eq.(1.3) for $\mu_1, \mu_2 \in \mathcal{M}_{\mathbb{R}_+}$, and therefore involves positive operators.

Theorem 7.1. *Let X_ι , $\iota \in I$, be positive random variables which are free with respect to φ , and denote $t_\iota = 1 + x_\iota$, $T_{\bar{\iota}} = \sqrt{X_{\bar{\iota}} - x_{\bar{\iota}}}X_{\bar{\iota}}\sqrt{X_{\bar{\iota}} - x_{\bar{\iota}}}$. Then $X_{\bar{\iota}}X_\iota$ has the same φ -distribution as $\sqrt{T_{\bar{\iota}}T_{\bar{\iota}}}\sqrt{t_\iota}$ and the pair $(t_\iota - 1, T_{\bar{\iota}} - 1)$ is monotone independent with respect to φ .*

Proof. Note that $T_{\bar{\iota}}$ introduced above is well-defined since

$$X_{\iota} - x_{\iota} = P_{\iota}(1) + \sum_{j=2}^{\infty} X_{\iota}(j)$$

is positive as an orthogonal direct sum of positive random variables. Using a similar argument, we get positivity of

$$t_{\iota} = X_{\iota}(1) + \sum_{j=2}^{\infty} P_{\iota}(j).$$

Computations of square roots give

$$\sqrt{t_{\iota}} = \sqrt{X_{\iota}(1)} + \sum_{j=2}^{\infty} P_{\iota}(j) \quad \text{and} \quad \sqrt{X_{\iota} - x_{\iota}} = P_{\iota}(1) + \sum_{j=2}^{\infty} \sqrt{X_{\iota}(j)},$$

where $\sqrt{X_{\iota}(j)} = P_{\iota}(j)\lambda(\sqrt{z})P_{\iota}(j)$ for $j \in \mathbb{N}$, with $z = x$ or $z = y$ if $\iota = 1$ or $\iota = 2$, respectively, which leads to

$$\sqrt{t_{\iota}}\sqrt{X_{\iota} - x_{\iota}} = \sqrt{X_{\iota}}.$$

Now, we use the fact that $X_{\bar{\iota}}X_{\iota}$ has the same φ -distribution as $\sqrt{X_{\iota}}X_{\bar{\iota}}\sqrt{X_{\iota}}$, which completes the proof of the first part of the theorem.

We will show now that the pair $(t_{\iota} - 1, T_{\bar{\iota}} - 1)$ is monotone independent w.r.t. φ . For simplicity, we denote $x = t_{\iota} - 1$ and $y = T_{\bar{\iota}} - 1$.

Case 1. If y^n is in the middle of the moment, we compute $\varphi(w_1xy^nxw_2)$, where $n \in \mathbb{N}$ and $w_1, w_2 \in \text{alg}(x, y)$. Note that the range of x is $\mathbb{C} \oplus \mathcal{H}_{\iota}^0$, therefore, we only need to find the action of y^n onto ξ and $h \in \mathcal{H}_{\iota}^0$. Using the explicit form of $\sqrt{X_{\iota} - x_{\iota}}$, we get

$$\begin{aligned} y^n \xi &= ((1 + \sum_{k=2}^{\infty} r_k)(1 + \sum_{j=1}^{\infty} z_j)(1 + \sum_{k=2}^{\infty} r_k) - 1)^n \xi \\ &= \varphi(y^n) \xi \quad \text{mod } \mathcal{K}(\iota), \end{aligned}$$

where $r_k = \sqrt{X_{\iota}(k)} - P_{\iota}(k)$ and $z_j = X_{\bar{\iota}}(j) - P_{\bar{\iota}}(j)$ and therefore $r_k \xi = 0$ for any $k \geq 2$. Similarly, we get

$$y^n h = \langle y^n h, h \rangle h \quad \text{mod } \mathcal{K}(\bar{\iota}) \ominus \mathcal{K}^{(1)}(\bar{\iota})$$

for any $h \in \mathcal{H}_{\iota}^0$ of norm $\|h\| = 1$. Note that h plays the role of a cyclic vector for the restriction of the free product representation $\pi_1 * \pi_2$ to the unital algebra generated by y . Thus $\langle y^n h, h \rangle = \varphi(y^n)h$, which gives

$$\varphi(w_1xy^nxw_2) = \varphi(y^n)\varphi(w_1x^2w_2),$$

i.e. the required condition for monotone independence.

Case 2. If y^n is at the end of the moment, we get

$$\varphi(w_1xy^n) = \varphi(y^n)\varphi(w_1x),$$

using the relation for $y^n \xi$ obtained above and the fact that $\mathcal{K}(\iota) \subset \text{Ker } x$, which finishes the proof. ■

Corollary 7.1. *If, in addition, X_{ι} has φ -distribution $\mu_{\iota} \neq \delta_0$ for $\iota \in I$, then the φ -distribution of T_{ι} is given by the s -free multiplicative convolution $\mu_{\iota} \boxplus \mu_{\bar{\iota}}$.*

Proof. First, note that the variable t_ι has φ -distribution μ_ι . In view of (1.5) and Theorem 7.1, we have $\eta_{\mu_1 \boxtimes \mu_2} = \eta_{\mu_1} \circ \eta_{\sigma_2} = \eta_{\mu_2} \circ \eta_{\sigma_1}$, where σ_1 and σ_2 are the φ -distributions of T_1 and T_2 , respectively. From the definition of the s-free multiplicative convolution, it follows that $\sigma_1 = \mu_1 \boxminus \mu_2$ and $\sigma_2 = \mu_2 \boxminus \mu_1$, which proves our assertion. ■

Remark 7.1. In analogy to the terminology used in analytic subordination [8,23], we can say that $X_{\bar{\tau}}X_\iota$ is ‘subordinate to t_ι and $t_{\bar{\tau}}$, with $T_{\bar{\tau}}$ and T_ι , respectively, being the corresponding ‘subordination operators’. Let us remark, however, that operators T_ι , $\iota \in I$, are not good candidates for ‘subordination branches’ of the product of free random variables since they are not suitable for further decomposition. In that sense, they are not multiplicative analogues of the ‘additive subordination branches’ [16], or the branches of the free product of graphs introduced by Quenell [20]. In the next section we will introduce and study such analogues, whose decompositions will correspond to decompositions of s-free multiplicative convolutions.

A similar operatorial subordination result can be established for bounded operators which includes unitary operators (the latter are related to $\mathcal{M}_{\mathbb{T}}$).

Theorem 7.2. *Let X_ι , $\iota \in I$, be bounded random variables which are free with respect to φ . Then, for each $\iota \in I$, the operator $X_{\bar{\tau}}X_\iota$ has the same φ -distribution as $Q_{\bar{\tau}}q_\iota$, where $q_\iota = 1 + x_\iota$ and $Q_{\bar{\tau}} = X_{\bar{\tau}}(X_\iota - x_\iota)$. Moreover, the pair $(q_\iota - 1, Q_{\bar{\tau}} - 1)$ is monotone independent with respect to φ . Finally, if X_ι , $\iota \in I$, are unitary, then the operators q_ι, Q_ι , $\iota \in I$, are unitary.*

Proof. The proof of the statements concerning bounded operators is similar to that of Theorem 7.1. Therefore, we shall just prove unitarity of Q_ι and q_ι . Writing operators q_ι and $X_\iota - x_\iota$ in the form of orthogonal series

$$q_\iota = X_\iota(1) + \sum_{j=2}^{\infty} P_\iota(j), \quad X_\iota - x_\iota = P_\iota(1) + \sum_{j=2}^{\infty} X_\iota(j),$$

we obtain

$$q_\iota q_\iota^* = X_\iota(1)X_\iota^*(1) + \sum_{j=2}^{\infty} P_\iota(j) = 1$$

and

$$(X_\iota - x_\iota)(X_\iota - x_\iota)^* = P_\iota(1) + \sum_{j=2}^{\infty} X_\iota(j)X_\iota^*(j) = 1.$$

Similarly, $q_\iota^* q_\iota = 1$ and $(X_\iota - x_\iota)^*(X_\iota - x_\iota) = 1$. This proves unitarity of q_ι and $X_\iota - x_\iota$, from which we obtain unitarity of Q_ι . The remaining part of the proof is similar to that of Theorem 7.1. ■

Corollary 7.2. *Let μ_ι and σ_ι be the φ -distributions of X_ι and Q_ι , respectively, for $\iota \in I$. Then the φ -distribution of $X_{\bar{\tau}}X_\iota$ is given by $\mu_\iota \cup \sigma_{\bar{\tau}}$ for each $\iota \in I$. In particular, if the operators X_ι are unitary, then $\sigma_\iota = \mu_\iota \boxminus \mu_{\bar{\tau}}$ for each $\iota \in I$.*

Proof. The first statement follows from Theorem 7.3 and the definition of the monotone multiplicative convolution of distributions. The proof of the second statement is

similar to that of Corollary 7.2 (the s-free convolution of measures from $\mathcal{M}_*$ is used). ■

8. SUBORDINATION BRANCHES

In order to introduce operator-valued ‘multiplicative subordination branches’ associated with the product of bounded free random variables, let us introduce random variables

$$(8.1) \quad R_\iota(m) = 1 + \sum_{j \in J(m)} x_\iota(j)$$

where the notation $J(m) = \{m, m+2, m+4, \dots\}$ is used for index sets and, by abuse of notation, 1 denotes $1_{\mathcal{K}_\iota}$ if m is odd, or $1_{\mathcal{K}_{\bar{\iota}}}$ if m is even. Note that operators $R_\iota(m)$ are elements of $B(\mathcal{K}_\iota)$ or $B(\mathcal{K}_{\bar{\iota}})$, depending on whether we have m odd or even, respectively.

Therefore, on each of the ‘s-free Fock spaces’, $\mathcal{K}_\iota$, $\iota \in I$, we get an ‘interlaced’ sequence of operators

$$(8.2) \quad R_\iota(1), R_{\bar{\iota}}(2), R_\iota(3), R_{\bar{\iota}}(4), \dots \in B(\mathcal{K}_\iota).$$

Each of these two sequences is used in the definition of *one* sequence of operator-valued ‘multiplicative subordination branches’.

Proposition 8.1. *If X_1, X_2 are positive (unitary), then operators $R_\iota(m)$, where $\iota \in I$ and $m \in \mathbb{N}$, are positive (unitary).*

Proof. This fact can be easily checked by decomposing the units according to (7.11). We choose to show unitarity of

$$R_\iota^{\text{odd}} = \sum_{j \text{ odd}} X_\iota(j) \quad \text{and} \quad R_\iota^{\text{even}} = P_\xi + \sum_{j \text{ even}} X_\iota(j)$$

for unitary X_ι . Since $X_\iota(j)X_\iota^*(j) = P_\iota(j)X_\iota X_\iota^* P_\iota(j) = P_\iota(j)$ for each $\iota \in I$ and $j \in \mathbb{N}$, we get

$$R_\iota^{\text{odd}}(R_\iota^{\text{odd}})^* = \sum_{j \text{ odd}} X_\iota(j)X_\iota^*(j) = \sum_{j \text{ odd}} P_\iota(j) = 1,$$

$$R_\iota^{\text{even}}(R_\iota^{\text{even}})^* = P_\xi + \sum_{j \text{ even}} X_\iota(j)X_\iota^*(j) = P_\xi + \sum_{j \text{ even}} P_\iota(j) = 1.$$

Unitarity of $R_\iota(m)$ for arbitrary m can be proved in similar way. Positivity of $R_\iota(m)$ follows easily from the appropriate decomposition of the unit. ■

Definition 8.1. Let X_ι , $\iota \in I$, be random variables which are free w.r.t. φ . Random variables

$$(8.3) \quad U_\iota(m) = R_\iota(m)R_{\bar{\iota}}(m+1),$$

where $\iota \in I$ and $m \in \mathbb{N}$, will be called *multiplicative bounded subordination branches*. If X_ι , $\iota \in I$ are unitary, then they will be called *multiplicative unitary subordination branches*.

In addition to branches $U_\iota(m)$, we shall need operators obtained from $U_\iota(m)$ by ‘order reversal’. Namely, let

$$(8.4) \quad V_\iota(m) = R_{\bar{\iota}}(m+1)R_\iota(m)$$

where $\iota \in I$ and $m \in \mathbb{N}$. Again, we use a simpler notation for $m = 1$, namely $V_\iota = V_\iota(m)$.

Definition 8.2. Let X_ι , $\iota \in I$ be positive random variables which are free w.r.t. φ . Random variables

$$(8.5) \quad Y_\iota(m) = \sqrt{R_{\bar{\iota}}(m+1)}R_\iota(m)\sqrt{R_{\bar{\iota}}(m+1)},$$

where $\iota \in I$ and $m \in \mathbb{N}$, will be called *multiplicative positive subordination branches*.

For simplicity, we will also use the terms: ‘bounded branches’, ‘unitary branches’ and ‘positive branches’. As in the case of operators $R_\iota(m)$, where $m \in \mathbb{N}$ and $\iota \in I$, we get two sequences of alternating (bounded, unitary) branches

$$(8.6) \quad U_\iota(1), U_{\bar{\iota}}(2), U_\iota(3), U_{\bar{\iota}}(4), \dots \in B(\mathcal{K}_\iota)$$

for each $\iota \in I$, and similar sequences of positive branches. Recall that existence of ‘interlaced branches’ was also observed for the additive (self-adjoint) subordination branches [16].

Of particular interest are branches of 1st order, and it is therefore of advantage to use a special notation for the first two operators in each ‘interlaced’ sequence given by (8.1). For $\iota \in I$, we shall use

$$(8.7) \quad R_\iota^{\text{odd}} = 1 + \sum_{j \text{ odd}} x_\iota(j) \quad \text{and} \quad R_\iota^{\text{even}} = 1 + \sum_{j \text{ even}} x_\iota(j)$$

where again, the unit 1 has to be interpreted as $1_{\mathcal{K}_\iota}$ and $1_{\mathcal{K}_{\bar{\iota}}}$, respectively. Note that although we have two different units here, the operators which appear in the same ‘interlaced’ sequence contain the same unit.

Remark 8.1. Let us recall that in the additive case [15, Theorem 8.4], the branches B_1 and B_2 of the sum $X_1 + X_2$ can be decomposed as

$$(8.8) \quad B_1 = S_1^{\text{odd}} + S_2^{\text{even}},$$

where $S_1^{\text{odd}} = \sum_{j \text{ odd}} X_1(j)$, $S_2^{\text{even}} = \sum_{j \text{ even}} X_2(j)$, and that the pair $(S_1^{\text{odd}}, S_2^{\text{even}})$ is s-free w.r.t. the pair of states (φ, ψ) , where ψ is associated with any unit vector $\zeta \in \mathcal{H}_1^0$. Note that we have $R_1^{\text{odd}} = S_1^{\text{odd}}$ and $R_2^{\text{even}} = P_\xi + S_2^{\text{even}}$, but we prefer to have a new notation to maintain a uniform style for ‘subtracting units’.

This, in turn, leads to a special notation for branches of 1st order, $U_\iota = U_\iota(1)$, $Y_\iota = Y_\iota(1)$, $\iota \in I$. For instance,

$$\begin{aligned} Y_1 &= Y_1(1) = \sqrt{R_2^{\text{even}}}R_1^{\text{odd}}\sqrt{R_2^{\text{even}}}, \\ Y_2 &= Y_2(1) = \sqrt{R_1^{\text{even}}}R_2^{\text{odd}}\sqrt{R_1^{\text{even}}}. \end{aligned}$$

Many computations can be reduced to branches of 1st order since their φ -distributions agree with the distributions of branches of higher orders with respect to suitably chosen states.

Let us examine the φ -distributions of branches of first order. In fact, we will show that Y_ι and U_ι have the same φ -distributions as T_ι for given $\iota \in I$.

Lemma 8.1. *The variables Q_ι , U_ι and V_ι have the same φ -distributions for any given $\iota \in I$. If X_ι , $\iota \in I$, are positive, then the variables $T_\iota, Y_\iota, U_\iota$ and V_ι have the same φ -distributions for any given $\iota \in I$.*

Proof. We shall give the proof in the more difficult case of positive operators (the proof for the case of bounded operators is similar). For $n \geq 1$, we have

$$\begin{aligned}\varphi(Y_\iota^n) &= \left(\sqrt{R_\iota^{\text{even}}} (R_\iota^{\text{odd}} R_\iota^{\text{even}})^{n-1} R_\iota^{\text{odd}} \sqrt{R_\iota^{\text{even}}} \right) \\ &= \varphi \left((R_\iota^{\text{odd}} R_\iota^{\text{even}})^{n-1} R_\iota^{\text{odd}} \right) \\ &= \varphi(U_\iota^n)\end{aligned}$$

since $\sqrt{R_\iota^{\text{even}}} \xi = \xi$ (the only summand in the direct sum decomposition of this square root which gives a non-zero contribution is $1_\tau(1)$). This shows that Y_ι and U_ι have the same φ -distributions.

Let us now compare the moments of T_ι with those of Y_ι . We have

$$\begin{aligned}\varphi(T_\iota^n) &= \varphi \left(\sqrt{X_\iota - x_\iota} (X_\iota (X_\iota - x_\iota))^{n-1} X_\iota \sqrt{X_\iota - x_\iota} \right) \\ &= \varphi((X_\iota (X_\iota - x_\iota))^n)\end{aligned}$$

since $\sqrt{X_\iota - x_\iota} \xi = \xi$. To fix attention, suppose that $\iota = 2$ and denote, for convenience, $z_k = x_1(k)$ and $y_k = x_2(k)$. Now, let us observe that in the finite set of variables of type z_k and y_j , where $k \geq 2$ and $j \geq 1$, which are used when calculating $X_2(X_1 - x_1)^{n-1} X_2 \xi$ there are no z_k 's for odd k 's, or y_j 's for even j 's. This is because the only variable which may give a non-zero contribution when acting on ξ is y_1 , then only y_1 and z_2 may follow, of which the first operator can be followed by y_1 or z_2 , whereas the second – by z_2, y_1 or y_3 , etc. This leads to

$$\varphi(P(X_2, X_1 - x_1)) = \varphi(P(R_2^{\text{odd}}, R_1^{\text{even}}))$$

for any polynomial P in two noncommuting variables. This shows that T_2 and Y_2 have the same φ -distributions.

Finally, we will show that U_2 and V_2 have the same φ -distributions. Using the notation used in the previous paragraph, we have

$$\begin{aligned}\varphi((U_2 - 1)^n) &= \varphi \left(\left(\sum_{k \text{ even}} z_k + \sum_{j \text{ odd}} y_j + \sum_{\substack{k \text{ even}, j \text{ odd} \\ |j-k|=1}} y_j z_k \right)^n \right) \\ \varphi((V_2 - 1)^n) &= \varphi \left(\left(\sum_{k \text{ even}} z_k + \sum_{j \text{ odd}} y_j + \sum_{\substack{k \text{ even}, j \text{ odd} \\ |j-k|=1}} z_k y_j \right)^n \right)\end{aligned}$$

for any natural n , and therefore, we need to show that the right-hand-sides of the above equations are equal to each other. Note that the products of variables z_k and y_j in words w which appear in mixed moments $\varphi(w)$ giving non-zero contributions to the above moments satisfy the following three conditions:

- (i) whenever z_k stands next to z_m or y_k next to y_m , it holds that $k = m$,
- (ii) whenever y_j stands next to z_k , it holds that $|j - k| = 1$,

(iii) the word w begins and ends with the letter y_1 .

Let us denote by W the set of words in letters

$$L = \{z_k, y_j : k \text{ even}, j \text{ odd}\}$$

subject to conditions (i)-(iii). In turn, let W' and W'' , respectively, denote the sets of all words in letters

$$\begin{aligned} L' &= \{z_k, y_j, z_k y_{k-1}, z_k y_{k+1} : k \text{ even}, j \text{ odd}\}, \\ L'' &= \{z_k, y_j, y_{k-1} z_k, y_{k+1} z_k : k \text{ even}, j \text{ odd}\} \end{aligned}$$

subject to conditions (i) and (iii), where products of type $z_k y_j$ and $y_j z_k$ are treated as letters. When they are used to form words, they are denoted by $[z_k y_j]$ and $[y_j z_k]$, respectively. Note that $W \subset W'$ and $W \subset W''$. More importantly, there is a bijection

$$\tau : W' \rightarrow W''$$

defined by ‘shifting brackets to the left’. Namely, τ is uniquely defined by $\tau(w_1) = w_1$ and the recursion

$$\tau(w_1 y_m z_k^p [z_k y_j] w_2) = w_1 [y_m z_k] z_k^p \tau(y_j w_2)$$

for any $w_1 \in W$, $w_2 \in W'$, $p \geq 0$ and $|j - k| = 1$, $|k - m| = 1$, i.e. we assume that $[z_k y_j]$ is the first such pair (counting from the left). In this recursion, we shift the brackets to the left from this pair to the closest possible pair y_m, z_k lying to its left (in view of (iii), such a pair must exist). It is not hard to see that its inverse is given by $\tau^{-1}(u_2) = u_2$ and the recursion

$$\tau^{-1}(w'_1 [y_j z_k] z_k^p y_m w'_2) = \tau^{-1}(w'_1 y_j) z_k^p [z_k y_m] w'_2$$

where $w'_1 \in W''$, $w'_2 \in W$, $p \geq 0$ and $|j - k| = 1$, $|k - m| = 1$. Again, note that for every such pair $[y_j z_k]$ there must exist y_m standing to its right in view of (iii). Clearly, for every $w \in W'$ it holds that $\varphi(\tau(w)) = \varphi(w)$. This completes the proof. $\blacksquare$

We are ready to prove a theorem, which will lead to a relation between the distributions of two consecutive branches. For that purpose, introduce operators

$$(8.9) \quad Z_\iota(m) = 1 + x_\iota(m)$$

with the same convention concerning the units as before ($1 = 1_{\kappa_\iota}$ if m is odd and $1 = 1_{\kappa_{\bar{\iota}}}$ if m is even), as well as operators obtained from the subordination branches by subtracting the (appropriate) unit:

$$(8.10) \quad y_\iota(m) = Y_\iota(m) - 1$$

for any $m \in \mathbb{N}$ and $\iota \in I$.

Theorem 8.1. *For given $m \in \mathbb{N}$ and $\iota \in I$, let φ, ψ be states associated with any unit vectors $\zeta \in \mathcal{H}^{(m-1)}(\iota)$ and $\zeta' \in \mathcal{H}^{(m)}(\bar{\iota})$. Then, operators $U_\iota(m)$ and $Z_\iota(m)U_{\bar{\iota}}(m+1)$ have the same φ -distributions. Similarly, if $X_\iota, \iota \in I$, are positive, then operators $Y_\iota(m)$ and $Z_\iota(m)Y_{\bar{\iota}}(m+1)$ have the same φ -distributions. Moreover, the pairs $(x_\iota(m), u_{\bar{\iota}}(m+1))$ and $(x_\iota(m), y_{\bar{\iota}}(m+1))$ are orthogonal with respect to (φ, ψ) for each $\iota \in I$ and $m \in \mathbb{N}$.*

Proof. We shall prove the more difficult case of positive random variables. Without loss of generality, let $\iota = 1$ and, for notational simplicity, consider the case $m = 1$ (the proof of the general case is similar). Denote

$$Z_1 = Z_1(1), R_1 = R_1(1), R_3 = R_1(3), R_2 = R_2(2), Y_2 = Y_2(2).$$

In the first part of the theorem we need to prove that Y_1 has the same φ -distribution as $Z_1 Y_2$. Using orthogonal decompositions and ‘square root calculus’, we obtain

$$Z_1 = X_1(1) + \sum_{j \in J(3)} P_1(j), \quad \text{and} \quad \sqrt{R_3} = P_1(1) + \sum_{j \in J(3)} \sqrt{X_1(j)}$$

which gives

$$\sqrt{R_3} Z_1 \sqrt{R_3} = \sum_{j \text{ odd}} X_1(j) = R_1,$$

and therefore

$$Y_1 = \sqrt{R_2} R_1 \sqrt{R_2} = \sqrt{R_2} \sqrt{R_3} Z_1 \sqrt{R_3} \sqrt{R_2}.$$

Therefore,

$$\begin{aligned} \varphi(Y_1^n) &= \varphi\left((\sqrt{R_2} \sqrt{R_3} Z_1 \sqrt{R_3} \sqrt{R_2})^n\right) \\ &= \varphi\left(\sqrt{R_2} \sqrt{R_3} (Z_1 \sqrt{R_3} R_2 \sqrt{R_3})^{n-1} Z_1 \sqrt{R_3} \sqrt{R_2}\right) \\ &= \varphi\left(\sqrt{R_2} \sqrt{R_3} (Z_1 Y_2)^{n-1} Z_1 \sqrt{R_3} \sqrt{R_2}\right) \\ &= \varphi\left((Z_1 Y_2)^{n-1} Z_1\right) \\ &= \varphi\left((Z_1 Y_2)^n\right) \end{aligned}$$

since $Y_2 \xi = \xi$ and also $\sqrt{R_3} \sqrt{R_2} \xi = \xi$. Therefore, the φ -distribution of Y_1 agrees with the φ -distribution of $Z_1 Y_2$.

In the proof of orthogonality, denote, as before, $x_1 = x_1(1)$, and $y_2 = Y_2 - 1_{\mathcal{K}_1}$. Using orthogonal decompositions and ‘square root calculus’, we can write

$$\begin{aligned} Y_2 &= \sqrt{R_3} R_2 \sqrt{R_3} \\ &= (P_1(1) + \sum_{j \in J(3)} \sqrt{X_1(j)})(P_\xi + \sum_{k \in J(2)} X_2(k))(P_1(1) + \sum_{j \in J(3)} \sqrt{X_1(j)}) \end{aligned}$$

and therefore $Y_2 \xi = \xi$, which implies that $y_2 \xi = 0$. Therefore, we get $\varphi(w y_2) = 0$ and therefore, by taking the adjoints, $\varphi(y_2 w) = 0$ for any $w \in \text{alg}(x_1, y_2)$. This gives the first orthogonality condition. Now, let us prove the second orthogonality condition, i.e.

$$\varphi(w_1 x_1 y_2^k x_1 w_2) = \psi(y_2^k)(\varphi(w_1 x_1^2 w_2) - \varphi(w_1 x_1) \varphi(x_1 w_2))$$

for any $k \in \mathbb{N}$ and $w_1, w_2 \in \text{alg}(x_1, y_2)$. Note that the ‘lowest order term’ in the expression for y_2 , which turns out to be of special importance when x_1 stands next to y_2 , is of the form

$$P_1(1) X_2(2) P_1(1) = P_1(1) P_2(2) X_2 P_2(2) P_1(1).$$

Recall that $P_2(2)$ is the projection onto $\mathcal{H}_1^0 \oplus (\mathcal{H}_2 \otimes \mathcal{H}_1^0)$ and $P_1(1)$ is the projection onto $\mathbb{C} \xi \oplus \mathcal{H}_1^0$, thus $P_1(1) P_2(2) = P_{\mathcal{H}_1^0}$, and therefore we get

$$x_1 y_2^m x_1 = x_1 P_{\mathcal{H}_1^0} y_2^m P_{\mathcal{H}_1^0} x_1,$$

for $m \geq 1$. This gives

$$\varphi(w_1 x_1 y_2^m x_1 w_2) = \varphi(w_1 x_1 P_{\mathcal{H}_1^0} x_1 w_2) \psi(y_2^m),$$

since, for any $\zeta \in \mathcal{H}_1^0$ of norm one, we have

$$y_2^m \zeta = \psi(y_2^m) \zeta \pmod{\mathcal{K}(2) \ominus \mathcal{K}^{(1)}(2)},$$

where ψ is the state associated with ζ . Note that ψ gives the same moments of the variable y_2 irrespective which ζ is taken and they agree with the corresponding moments in the state φ_2 . Finally, since $P_{\mathcal{H}_1^0} = P_{\mathbb{C}\xi \oplus \mathcal{H}_1^0} - P_{\mathbb{C}\xi}$, we have

$$\varphi(w_1 x_1 P_{\mathcal{H}_1^0} x_1 w_2) = \varphi(w_1 x_1^2 w_2) - \varphi(w_1 x_1) \varphi(x_1 w_2),$$

which completes the proof of the second orthogonality condition. $\blacksquare$

Corollary 8.1. *Let μ_ι and σ_ι be φ -distributions of (bounded operators) X_ι and U_ι , respectively, for each $\iota \in I$, and let ψ_{m-1} be the state associated with any unit vector $\zeta \in \mathcal{H}^{(m-1)}(\iota)$, where $m \in \mathbb{N}$. Then*

- (1) *the ψ_{m-1} -distribution of $U_\iota(m)$ agrees with σ_ι for every $m \geq 2$ and $\iota \in I$,*
- (2) *it holds that $\sigma_\iota = \mu_\iota \angle \sigma_\tau$ for each $\iota \in I$,*
- (3) *if X_1, X_2 are positive and $\mu_1, \mu_2 \neq \delta_0$, then σ_ι agrees with the s -free convolution $\mu_\iota \boxtimes \mu_\tau$ of probability measures on $\mathbb{R}_+$ for each $\iota \in I$,*
- (4) *if X_1, X_2 are unitary, then σ_ι agrees with the s -free convolution $\mu_\iota \boxtimes \mu_\tau$ of probability measures on $\mathbb{T}$ for each $\iota \in I$.*

Proof. Assertion (1) follows from the definition of operators $U_\iota(m)$. Then, (2) is a consequence of (1) and Theorem 8.3. Finally, it follows from Theorems 7.3 and 8.3 that the φ -distributions of U_ι satisfy the subordination equations (2.1). Uniqueness of the subordination functions for the cases considered in (3) and (4) give the assertions. $\blacksquare$

Corollary 8.1, together with Corollary 4.6, naturally lead to the definition of a sequence of iterations of the s -free multiplicative convolution. Thus, for given $\mu_1, \mu_2 \in \Sigma$ let

$$(8.11) \quad \mu_1 \angle_1 \mu_2 = \mu_1 \angle \mu_2, \quad \mu_1 \angle_n \mu_2 = \mu_1 \angle (\mu_2 \angle_{n-1} \mu_1),$$

for $n \geq 2$ (of course, this sequence can also be defined by for any measures, for which the operation $\angle$ has been defined).

Corollary 8.2. *If $\mu_1, \mu_2 \in \mathcal{M}_{\mathbb{R}_+} \setminus \{\delta_0\}$ are compactly supported, then*

$$(8.12) \quad w - \lim_{n \rightarrow \infty} (\mu_1 \angle_n \mu_2) = \mu_1 \boxtimes \mu_2$$

$$(8.13) \quad w - \lim_{n \rightarrow \infty} (\mu_1 \circ (\mu_2 \angle_n \mu_1)) = \mu_1 \boxtimes \mu_2,$$

and the corresponding η -transforms converge uniformly on the compact subsets of $\mathbb{C} \setminus \mathbb{R}_+$ to the η -transforms of the limit measures. An analogous result holds for $\mu_1, \mu_2 \in \mathcal{M}_*$.

Proof. In view of Corollary 4.6, for fixed $m \in \mathbb{N}$, $(\mu_1 \angle_n \mu_2)(m) = (\mu_1 \angle_m \mu_2)(m)$ for all $n \geq m$, from which we obtain convergence of moments as $n \rightarrow \infty$ for any distributions $\mu_1, \mu_2 \in \Sigma$. Since the measures are compactly supported, this implies weak convergence

of the corresponding measures. Therefore, the η -transforms converge uniformly on compact subsets of $\mathbb{C} \setminus \mathbb{R}_+$. Moreover, by Corollary 8.1, the limit distributions agree with the distributions of the positive subordination branch Y_1 , which is $\mu_1 \boxtimes \mu_2$. The arguments for the weak convergence of the second sequence are similar. An analogous proof holds for $\mu_1, \mu_2 \in \mathcal{M}_*$. $\blacksquare$

Remark 8.2. Informally, the weak limits of Corollary 8.2 can be written in the following form:

$$\begin{aligned}\mu_1 \boxtimes \mu_2 &= \mu_1 \angle (\mu_2 \angle (\mu_1 \angle (\mu_2 \angle (\dots))))), \\ \mu_1 \boxtimes \mu_2 &= \mu_1 \circ (\mu_2 \angle (\mu_1 \angle (\mu_2 \angle (\dots))))),\end{aligned}$$

whereas their transforms in the ‘continued composition form’:

$$\begin{aligned}\rho_{\mu_1 \boxtimes \mu_2}(z) &= \rho_{\mu_1}(z \rho_{\mu_2}(z \rho_{\mu_1}(z \rho_{\mu_2}(\dots)))), \\ \eta_{\mu_1 \boxtimes \mu_2}(z) &= \eta_{\mu_1}(z \rho_{\mu_2}(z \rho_{\mu_1}(z \rho_{\mu_2}(\dots)))),\end{aligned}$$

where the right-hand sides are understood as the uniform limits on compact subsets of $\mathbb{C} \setminus \mathbb{R}_+$ or $\mathbb{D}$. Actually, these formulas can be used to compute some examples, including simple examples of $\mu_1 \boxtimes \mu_2$ (without using S-transforms).

Remark 8.3. By Corollary 8.2, one can *define* the s-free and free convolutions of any compactly supported measures μ_1, μ_2 on $\mathbb{C}$ as the weak limits of the form (8.12)-(8.13). This allows us to use operatorial subordination results of Sections 7-8, including those which concern distributions of products of free bounded random variables as well as distributions of bounded branches and express them in terms of s-free convolutions, denoted with the same symbol $\boxtimes$ and understood as weak limits of type (8.12)-(8.13). Using the weak limits, we also get

$$(8.14) \quad \delta_0 \boxtimes \mu = \delta_0 \quad \text{and} \quad \mu \boxtimes \delta_0 = \delta_{\mu(X)}$$

and

$$(8.15) \quad \mu \boxtimes \delta_0 = \delta_0 = \delta_0 \boxtimes \mu,$$

for any compactly supported probability measure on $\mathbb{C}$, where we used the well-known relations $\mu \circ \delta_0 = \delta_0 = \delta_0 \circ \mu$.

Example 8.1. Using Corollary 8.2 and the results of Example 5.1, we obtain $\mu \boxtimes \delta_a = \mu \angle \delta_a = S_a(\mu)$ and $\delta_a \boxtimes \mu = \delta_a \angle \mu = \delta_a$ for compactly supported $\mu \in \mathcal{M}_{\mathbb{R}_+}$ and $a > 0$. This gives

$$\delta_a \boxtimes \mu = \delta_a \circ S_a \mu = D_a \mu,$$

in view of (1.5), since the corresponding η -transform is of the form

$$\eta_{\delta_a}(\eta_{S_a \mu}(z)) = \eta_{D_a \mu}(z),$$

where we used $\eta_{\delta_a}(z) = az$ and $\eta_{S_a \mu}(z) = \eta_{D_a \mu}(z)/a$. On the other hand,

$$\mu \boxtimes \delta_a = \mu \circ \delta_a = D_a \mu,$$

since $\eta_{\mu \circ \delta_a}(z) = \eta_{\mu}(az) = \eta_{D_a \mu}(z)$. In particular, we have $\delta_a \boxtimes \delta_b = \delta_a$ for $a, b > 0$.

Example 8.2. Using the results of Example 5.3, we obtain

$$\mu \boxtimes \delta_a = S_a \mu \quad \text{and} \quad \delta_a \boxtimes \mu = \delta_a,$$

for compactly supported $\mu \in \mathcal{M}_{\mathbb{T}}$ and $a \in \mathbb{T}$, which gives

$$\mu \boxtimes \delta_a = D_a \mu = \delta_a \boxtimes \mu,$$

by the same arguments as those used in Example 8.1.

Let us finally show that the multiplicative subordination branches are related to the notion of s-free independence, as it was the case of their additive counterparts. For that purpose, let us recall this concept [16].

Definition 8.3. Let $(\mathcal{A}, \varphi, \psi)$ be a unital algebra with a pair of linear normalized functionals. Let $\mathcal{A}_1$ be a unital subalgebra of $\mathcal{A}$ and let $\mathcal{A}_2$ be a non-unital subalgebra with an ‘internal’ unit 1_2 , i.e. $1_2 b = b = b 1_2$ for every $b \in \mathcal{A}_2$. We say that the pair $(\mathcal{A}_1, \mathcal{A}_2)$ is *free with subordination*, or simply *s-free*, with respect to (φ, ψ) if $\psi(1_2) = 1$ and it holds that

- (i) $\varphi(a_1 a_2 \dots a_n) = 0$ whenever $a_j \in \mathcal{A}_{i_j}^0$ and $i_1 \neq i_2 \neq \dots \neq i_n$
- (ii) $\varphi(w_1 1_2 w_2) = \varphi(w_1 w_2) - \varphi(w_1) \varphi(w_2)$ for any $w_1, w_2 \in \text{alg}(\mathcal{A}_1, \mathcal{A}_2)$,

where $\mathcal{A}_1^0 = \mathcal{A}_1 \cap \ker \varphi$ and $\mathcal{A}_2^0 = \mathcal{A}_2 \cap \ker \psi$. We say that the pair (a, b) of random variables from $\mathcal{A}$ is *s-free* with respect to (φ, ψ) if there exists $1_2 \in \mathcal{A}$ such that $(\mathcal{A}_1, \mathcal{A}_2)$, where $\mathcal{A}_1 := \text{alg}(1, a)$ and $\mathcal{A}_2 := \text{alg}(1_2, b)$, is s-free w.r.t. (φ, ψ) .

Proposition 8.2. *The pair $(R_\iota^{\text{odd}} - 1, R_\iota^{\text{even}} - 1)$ is s-free with respect to the pair (φ, ψ) , where ψ is the state associated with any unit vector $\zeta \in \mathcal{H}_\iota^0$, and the projection associated with the second variable is $1_2 = 1 - P_\xi$.*

Proof. For simplicity, choose $\iota = 1$ and denote $r_1 = R_1^{\text{odd}} - 1$, $r_2 = R_2^{\text{even}} - 1$. We have

$$r_1 = \sum_{j \text{ odd}} x_1(j) \quad \text{and} \quad r_2 = \sum_{j \text{ even}} x_2(j).$$

Clearly, the definition of 1_2 immediately gives $\psi(1_2) = 1$ as well as condition (ii) of s-freeness. Let us show that condition (i) also holds. Let $\mathcal{A}_1 = \text{alg}(r_1, 1)$ and $\mathcal{A}_2 = \text{alg}(r_2, 1_2)$. If in (i) we take $a_n \in \mathcal{A}_2 \cap \ker \psi$, then $\lambda(a_n) \xi = 0$, where λ is the free product representation, and hence (i) holds. If $a_n, a_{n-2}, \dots \in \mathcal{A}_1 \cap \ker \varphi$ and $a_{n-1}, a_{n-3}, \dots \in \mathcal{A}_2 \cap \ker \psi$, then the GNS representation space for such a mixed moment is $\mathcal{K}_1$ (none of $a_{n-1}, a_{n-3}, \dots$ ever gets to act on ξ). Therefore, in the computation of this kind of moment, one can replace r_1, r_2 and 1_2 by X_1, X_2 and 1, respectively. But then (i) follows from the usual freeness condition. $\blacksquare$

9. FREE PRODUCT AND S-FREE LOOP PRODUCT OF GRAPHS

In this section we prove the Multiplication Theorem for s-free independence and free independence.

Let us recall the definition of the s-free product of rooted graphs [2,16].




FIGURE 4. An example of the s-free loop product

Definition 9.1. By the *s-free* product of rooted graphs $(\mathcal{G}_1, e_1)$ and $(\mathcal{G}_2, e_2)$, denoted $(\mathcal{G}_1 \oplus \mathcal{G}_2, e_1)$, we understand the inductive limit of the sequence $(\mathcal{B}_1(m), e_1)_{m \in \mathbb{N}}$ of rooted graphs, where $\mathcal{B}_1(1) = \mathcal{G}_1 \vdash \mathcal{G}_2$ and $\mathcal{B}_l(m)$ is obtained from $\mathcal{B}_l(m-1)$ by attaching by its root a copy of $\mathcal{G}_1$ (if m is even), or a copy of $\mathcal{G}_2$ (if m is odd) to every vertex of the difference $\mathcal{B}_l(m-1) \setminus \mathcal{B}_l(m-2)$.

Remark 9.1. It can be seen that $(\mathcal{B}_1(m), e_1)_{m \in \mathbb{N}}$ is a sequence of growing graphs with the root kept to be e_1 , and therefore, the inductive limit exists. In a similar way we define $(\mathcal{G}_2 \oplus \mathcal{G}_1, e_2)$. It is easy to see that

$$(9.1) \quad \mathcal{B}_1(m) = \mathcal{G}_1 \vdash \mathcal{B}_2(m-1) \quad \text{and} \quad \mathcal{B}_2(m) = \mathcal{G}_2 \vdash \mathcal{B}_1(m-1),$$

which can also serve an inductive definition of the considered sequences. These formulas are important since they correspond to the decomposition of the s-free additive convolution of the form

$$(9.2) \quad \mu_1 \boxplus \mu_2 = \mu_1 \vdash (\mu_2 \vdash (\mu_1 \vdash (\mu_2 \vdash \dots))),$$

where μ_l is the spectral distribution of $\mathcal{G}_l$ [2].

Using the decomposition (9.2), we proved the Addition Theorem for s-free independence in [2]. Our first goal in this Section is to prove the corresponding Multiplication Theorem. For that purpose, we introduce a new notion of graph product, called the ‘s-free loop product’. We follow it up with a proposition which justifies the definition.

Definition 9.2. Suppose that the s-free product of graphs $(\mathcal{G}_1 \oplus \mathcal{G}_2, e_1)$ is naturally colored. The *s-free loop product* of $(\mathcal{G}_1, e_1)$ and $(\mathcal{G}_2, e_2)$ is the graph $(\mathcal{G}_1 \oplus_l \mathcal{G}_2, e_1)$ obtained from $(\mathcal{G}_1 \oplus \mathcal{G}_2, e_1)$ by attaching a loop of color 2 to the root e_1 .

Example 9.1. Consider the s-free loop product of graphs given in Fig.4, which can be viewed as a ‘binary tree with loops’ (the product graph is of course infinite, only our picture is truncated at distance 5 from the root). Note that graph $\mathcal{G}_1$ has a loop at the root, and so do copies of $\mathcal{G}_1$ in the product graph, whereas $\mathcal{G}_2$ does not have any loops. We produce $\mathcal{G}_1 \oplus \mathcal{G}_2$ in the inductive way (Definition 9.1 is used). Now, in order to obtain the s-free loop product, we attach one additional loop of color 2 to the root (we draw this loop larger than loops of color 1). On the level of the adjacency matrices, this additional loop corresponds to the projection P_ξ as the proof of Proposition 9.1 shows. Note that the main purpose of attaching this extra loop is to fit the s-free multiplicative convolution into the general scheme of the Multiplication Theorem, where counting

rooted alternating d-walks leads to the moments of multiplicative convolutions. Direct computations give, for instance, $D_2(e_1) = 1$, $D_4(e_1) = 0$, $D_6(e_1) = 4$, $D_8(e_1) = 0$.

Proposition 9.1. *Let A_ι be the adjacency matrix of $\mathcal{G}_\iota$, where $\iota \in I$. The adjacency matrix of the s-free loop product of $\mathcal{G}_1$ and $\mathcal{G}_2$ takes the form*

$$(9.3) \quad A(\mathcal{G}_1 \oplus_\ell \mathcal{G}_2) = R_1 + R_2$$

where

$$(9.4) \quad R_1 = \sum_{j \text{ odd}} A_1(j) \quad \text{and} \quad R_2 = P_\xi + \sum_{k \text{ even}} A_2(k).$$

Moreover, the φ_e -distribution of R_1 is μ_1 , the ψ -distribution of R_2 is μ_2 and the pair $(R_1 - 1, R_2 - 1)$ is s-free independent w.r.t. (φ_e, ψ) , where ψ is the state associated with any unit vector $\zeta \in \mathcal{H}_1^0 = l_2(V_1^0)$.

Proof. It follows from [2,16] that the adjacency matrix of the s-free product of graphs is of the form

$$A(\mathcal{G}_1 \oplus \mathcal{G}_2) = S_1 + S_2,$$

where

$$S_1 = \sum_{j \text{ odd}} A_1(j) \quad \text{and} \quad S_2 = \sum_{k \text{ even}} A_2(k).$$

By Definition 9.1, the adjacency matrix of the corresponding s-free loop product is obtained by adding the projection P_0 to the adjacency matrix of the subgraph of color 2, namely S_2 , which corresponds to attaching the loop of color 2 to the root. This gives

$$A(\mathcal{G}_1 \oplus_\ell \mathcal{G}_2) = R_1 + R_2,$$

where

$$R_1 = S_1 \quad \text{and} \quad R_2 = P_\xi + S_2.$$

In view of Proposition 8.2, the pair $(R_1 - 1, R_2 - 1)$ is s-free independent w.r.t. the pair (φ, ψ) , where ψ is the state associated with any unit vector $\zeta \in \mathcal{H}_1^0$ (here, 1 denotes the unit on $l_2(V)$, where V is the set of vertices of $\mathcal{G}_1 \oplus_\ell \mathcal{G}_2$, which corresponds to $\mathcal{K}_1$). Moreover, it is easy to check that the φ_e -distribution of R_1 is μ_1 and the ψ -distribution of R_2 is μ_2 . This completes the proof. $\blacksquare$

Thus, the first part of the Multiplication Theorem for s-free independence is proved. We now need to find a connection between the ‘first return moments’ of $R_2 R_1$ and cardinalities of the sets D_{2n} . In that context, note that in order that the set D of all rooted alternating d-walks on $\mathcal{G}_1 \oplus_\ell \mathcal{G}_2$ be non-empty, either $\mathcal{G}_1$ or $\mathcal{G}_2$ must have a loop at e_1 or e_2 , respectively. In fact, otherwise, starting an alternating walk from the root of $\mathcal{G}_1 \oplus_\ell \mathcal{G}_2$, one cannot return to it since the m -th edge of that walk must be of different color than the $m - 1$ -th edge and thus it must belong to $\mathcal{B}_j(m) \setminus \mathcal{B}_j(m - 1)$, and therefore its distance from the root (understood as the distance of the vertex closer to the root) equals $m - 1$ and thus tends to ∞ as $m \rightarrow \infty$.

Theorem 9.1. *The Multiplication Theorem holds for s-free independence, the associated loop product $\mathcal{G}_1 \oplus_\ell \mathcal{G}_2$, and the multiplicative convolution $\mu_1 \boxtimes \mu_2$.*

Proof. For notational simplicity, denote $\sigma_1 = \mu_1 \boxtimes \mu_2$, $\sigma_2 = \mu_2 \boxtimes \mu_1$ and

$$\mathcal{B}_\ell = \mathcal{G}_\ell \oplus \mathcal{G}_{\bar{\ell}}, \quad \mathcal{B}_\ell^\ell = \mathcal{G}_\ell \oplus_\ell \mathcal{G}_{\bar{\ell}}$$

From Proposition 9.1 and the results of Section 8, it follows that

$$(9.5) \quad N_{R_2 R_1}(n) = N_{\sigma_1}(n)$$

for $n \in \mathbb{N}$. Now, we need to prove that

$$(9.6) \quad N_{\sigma_1}(n) = |D_{2n}(e)|$$

for $k \in \mathbb{N}$. Note first that any f-walk $w \in F(e_1)$ on $\mathcal{B}_1$ must begin and terminate with an edge of color 1 since no edge of color 2 is incident on e_1 . Since $\mathcal{B}_1^\ell$ is obtained from $\mathcal{B}_1$ by attaching a loop of color 2 to the root e_1 , the set of rooted alternating d-walks on $\mathcal{B}_1^\ell$ of length $2n$ is in 1-1 correspondence with the set of rooted alternating f-walks on $\mathcal{B}_1$ of length $2n - 1$. In fact, each rooted alternating d-walk of length $2n$ on $\mathcal{B}_1^\ell$ is of the form $w = (u_1, u_2)$, where u_1 is a rooted alternating f-walk of length $2n - 1$ and u_2 is the rooted loop of color 2. Therefore,

$$(9.7) \quad |D_{2n}(e_1)| = |F_{2n-1}(e_1)|,$$

where $D_{2n}(e_1)$ refers to the loop product and $F_{2n-1}(e_1)$ – to the usual product. This fact will be used in the induction proof given below. If $n = 1$, we use this correspondence to get $N_{\sigma_1}(1) = N_{\mu_1}(1) = |F_1(e_1)| = |D_2(e_1)|$. Suppose now that (9.6) holds if n is replaced by $1 \leq k \leq n - 1$, where σ_1 is understood to be the s-free convolution of any σ' and σ'' (thus, in particular, μ_2 and μ_1). Using the proof of Theorem 6.1, we have

$$N_{\sigma_1}(n) = \sum_{r=1}^n N_{\mu_1}(r) \sum_{k_1+k_2+\dots+k_{r-1}=n-1} N_{\sigma_2}(k_1)N_{\sigma_2}(k_2)\dots N_{\sigma_2}(k_{r-1}).$$

In view of (9.7), it is enough to show that the expression on the RHS is equal to $|F_{2n-1}(e_1)|$. By the inductive assumption, we replace in the above formula each $N_{\sigma_2}(k_i)$ by $|D_{2k_i}(e_1)|$, and then, using (9.7), by $|F_{2k_i-1}(e_1)|$, $1 \leq i \leq r - 1$. Therefore, we need to justify the above formula, viewing $N_{\sigma_1}(n)$ as the number of rooted alternating f-walks on $\mathcal{B}_1$ and $N_{\sigma_2}(k_i)$, $1 \leq i \leq r - 1$, as numbers of rooted alternating f-walks on $\mathcal{B}_2$. Since

$$\mathcal{B}_1 = \mathcal{G}_1 \vdash \mathcal{B}_2$$

we can observe that each rooted alternating f-walk of length $2n - 1$ on $\mathcal{B}_1$ consists of a rooted f-walk $c = (v_0, v_1, \dots, v_r) \in F_r(e_1)$ of color 1, for some $1 \leq r \leq n$, with $r - 1$ alternating f-walks $w_i \in F(v_i)$ on copies of $\mathcal{B}_2$ attached to vertices v_i , where $1 \leq i \leq r - 1$. The latter must begin and terminate with edges of color 2 since no edge of color 1 is incident on the roots of these copies. Thus, the expression given on the RHS of the above formula gives the numbers of all alternating rooted f-walks of total length

$$r + (2k_1 - 1) + (2k_2 - 1) + \dots + (2k_{r-1} - 1) = r + 2(n - 1) - (r - 1) = 2n - 1$$

summed over $1 \leq r \leq n$, which completes the proof. ■

Example 9.2. The lowest order first return moments $N_{\mu_1 \boxtimes \mu_2}(n)$ are given by

$$\begin{aligned} N_{\mu_1 \boxtimes \mu_2}(1) &= N_{\mu_1}(1), \\ N_{\mu_1 \boxtimes \mu_2}(2) &= N_{\mu_1}(2)N_{\mu_2}(1), \end{aligned}$$

$$\begin{aligned}
 N_{\mu_1 \boxtimes \mu_2}(3) &= N_{\mu_1}(3)N_{\mu_2}^2(1) + N_{\mu_1}(2)N_{\mu_1}(1)N_{\mu_2}(2), \\
 N_{\mu_1 \boxtimes \mu_2}(4) &= N_{\mu_1}(4)N_{\mu_2}^3(1) + 2N_{\mu_1}(3)N_{\mu_1}(1)N_{\mu_2}(2)N_{\mu_2}(1) \\
 &\quad + N_{\mu_1}(2)N_{\mu_1}^2(1)N_{\mu_2}(3) + N_{\mu_1}^2(2)N_{\mu_2}(2)N_{\mu_2}(1)
 \end{aligned}$$

where we used the results of Example 6.2. The enumeration of d-walks given in Example 9.1 can be easily verified by substituting to the above formulas only the non-vanishing moments: $N_{\mu_1}(1) = 1$, $N_{\mu_1}(2) = 2$, $N_{\mu_2}(2) = 2$. We have $N_{\sigma_1}(1) = 1 = D_2(e_1)$, $N_{\sigma_1}(2) = 0 = D_4(e_1)$, $N_{\sigma_1}(3) = 4 = D_6(e_1)$ and $N_{\sigma_1}(4) = 0 = D_8(e_1)$, where $\sigma_1 = \mu_1 \boxtimes \mu_2$.

Finally, we will prove the Multiplication Theorem for free independence. In this case, one does not need to introduce a new type of graph product since no ‘unitization’ of adjacency matrices is necessary to make them freely independent (basically, this is because units are identified in the case of the free product of algebras). Nevertheless, if neither of the graphs, $\mathcal{G}_1$ or $\mathcal{G}_2$, have loops at their roots, then the set of rooted alternating d-walks on $\mathcal{G}_1 * \mathcal{G}_2$ is empty.

Theorem 9.2. *The Multiplication Theorem holds for free independence, the associated product graph $\mathcal{G}_1 * \mathcal{G}_2$, and the multiplicative convolution $\mu_1 \boxtimes \mu_2$.*

Proof. First of all, we know [2] that

$$A(\mathcal{G}_1 * \mathcal{G}_2) = S_1 + S_2$$

where $S_\iota = \sum_{k=1}^{\infty} A_\iota(k)$, $\iota \in I$. Moreover, the pairs (S_1, S_2) and $(S_1 - 1, S_2 - 1)$ are free w.r.t. φ . Also, it is clear that the moments of $S_2 S_1$ agree with the moments of $\mu_1 \boxtimes \mu_2$. It remains to be shown that the latter coincide with the corresponding numbers of rooted alternating d-walks. Using Proposition 2.1, we have $\mu_1 \boxtimes \mu_2 = \mu_1 \cup (\mu_2 \boxtimes \mu_1)$ and therefore, we can employ the combinatorial formula used in the proof of Theorem 3.1 to get

$$N_{\sigma_1}(n) = \sum_{r=1}^n N_{\mu_1}(r) \sum_{k_1+k_2+\dots+k_r=n} N_{\sigma_2}(k_1)N_{\sigma_2}(k_2)\dots N_{\sigma_2}(k_r).$$

Now, we can decompose the free product of graphs as

$$\mathcal{G}_1 * \mathcal{G}_2 = \mathcal{G}_1 \triangleright \mathcal{B}_2$$

with the root e of $\mathcal{G}_1 * \mathcal{G}_2$ obtained by identifying the root e_1 of $\mathcal{G}_1$ with the root e_2 of $\mathcal{B}_2$ (see [2,16,20]). In view of Proposition 9.1,

$$N_{\sigma_2}(k_i) = |F_{2k_i-1}(e_2)|,$$

where e_2 is the root in $\mathcal{B}_2$. Of course, $N_{\mu_1}(r) = |F_r(e_1)|$, where e_1 is treated as the root of $\mathcal{G}_1$. Therefore, the RHS of the above combinatorial formula gives the number of rooted walks w on $\mathcal{G}_1 * \mathcal{G}_2$ consisting of an f-walk $c = (v_0, v_1, \dots, v_r) \in F_r(e)$, interlaced with rooted alternating f-walks $w_1, w_2, \dots, w_r$ on copies of $\mathcal{B}_2$ attached to vertices $v_1, v_2, \dots, v_r$, respectively. In other words, we have

$$w = (\beta_1, w_1, \beta_2, w_2, \dots, \beta_r, w_r),$$




FIGURE 5. An example of the free product

and therefore, it consists of two rooted alternating f-walks, namely

$$u_1 = (\beta_1, w_1, \beta_2, w_2, \dots, \beta_r) \quad \text{and} \quad u_2 = w_r,$$

where u_1 begins and terminates with an edge of color 1, and u_2 begins and terminates with an edge of color 2. Therefore, w is a rooted alternating d-walk of length $2n$ which is ‘subordinate’ to an f-walk c of length r . The summation over $1 \leq r \leq n$ gives all rooted d-walks of length $2n$, which completes the proof. ■

In view of this result, the free product of graphs is ‘complete’ in our ‘category’ of product graphs since it is naturally related to the rooted alternating d-walks and no additional loops are necessary to fit it into the scheme of the Multiplication Theorem.

Example 9.3. The first return moments $N_{\mu_1 \boxtimes \mu_2}(n) = N_{\mu_2 \boxtimes \mu_1}(n)$ of lowest orders can be computed using (2.1) and the results of Example 3.2, with the $N_{\sigma_1}(n)$, where $\sigma_1 = \mu_1 \boxtimes \mu_2$, taken from Example 9.2. For the free product in Fig.5, we obtain $D_2(e) = 0$, $D_4(e) = 2$, $D_6(e) = 0$ and $D_8(e) = 16$.

REFERENCES

- [1] L. Accardi, A. Ben Ghorbal, N. Obata, Monotone independence, comb graphs and Bose-Einstein condensation, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **7** (2004), 419-435.
- [2] L. Accardi, R. Lenczewski, R. Śaławata, Decompositions of the free product of graphs, arXiv:math.CO/0609329, 2006.
- [3] S. Belinschi, Complex Analysis Methods in Noncommutative Probability, Ph.D. Thesis, Indiana University, 2005.
- [4] S. Belinschi, H. Bercovici, Partially defined semigroups relative to multiplicative free convolution, *Int. Math. Res. Not.* **2005**, no.2, 65-101.
- [5] H. Bercovici, On boolean convolutions, preprint, 2004.
- [6] H. Bercovici, Multiplicative monotonic convolution, *Illinois J. Math.* **49** (2005), 929-951.
- [7] H. Bercovici, D. Voiculescu, Free convolution of measures with unbounded support, *Indiana Univ. Math. J.* **42** (1993), 733-773.
- [8] Ph. Biane, Processes with free increments, *Math. Z.* **227** (1998), 143-174.
- [9] M. Bożejko, J. Wysocański, Remarks on t -transformations of measures and convolutions, *Ann.I.H.Poincaré- PR* **37**, 6 (2001), 737-761.

- [10] G.P. Chistyakov, F. Goetze, The arithmetic of distributions in free probability theory, preprint, arXiv:math.QA/0508245, 2005.
- [11] U. Franz, Boolean convolution of probability measures on the unit circle, preprint, 2004.
- [12] E. Gutkin, Green's functions of free products of operators with applications to graph spectra and to random walks, *Nagoya Math. J.* **149** (1998), 93-116.
- [13] R. Lenczewski, Unification of independence in quantum probability, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, **1** (1998), 383-405.
- [14] R. Lenczewski, Reduction of free independence to tensor independence, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **7** (2004), 337-360.
- [15] R. Lenczewski, R. Śaławata, Discrete interpolation between monotone probability and free probability, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **9** (2006), 77-106.
- [16] R. Lenczewski, Decompositions of the free additive convolution, *J. Funct. Anal.* (2007), doi:10.1016/j.jfa.2007.01.010.
- [17] H. Maassen, Addition of freely independent random variables, *J. Funct. Anal.* **106** (1992), 409-438.
- [18] N. Muraki, Monotonic independence, monotonic central limit theorem and monotonic law of small numbers, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **4** (2001), 39-58.
- [19] N. Muraki, Monotonic convolution and monotonic Levy-Hincin formula, preprint, 2001.
- [20] G. Quenell, Combinatorics of free product graphs, *Contemp. Math.* **206** (1994), 257-281.
- [21] D. Voiculescu, Symmetries of some reduced free product C^* -algebras, *Operator Algebras and their Connections with Topology and Ergodic Theory, Lecture Notes in Math.* 1132, Springer, Berlin, 1985, 556-588.
- [22] D. Voiculescu, Addition of certain non-commuting random variables, *J. Funct. Anal.* **66** (1986), 323-246.
- [23] D. Voiculescu, Multiplication of certain non-commuting random variables, *J. Operator Theory* **18** (1987), 223-235.
- [24] D. Voiculescu, The analogues of entropy and of Fisher's information measure in free probability theory, I, *Commun. Math. Phys.* **155** (1993), 71-92.
- [25] D. Voiculescu, K. Dykema, A. Nica, *Free random variables*, CRM Monograph Series, No.1, A.M.S., Providence, 1992.
- [26] W. Woess, *Random walks on Infinite Graphs and Groups*, Cambridge University Press, Cambridge, 2000.

ROMUALD LENCZEWSKI,
 INSTYTUT MATEMATYKI I INFORMATYKI, POLITECHNIKA WROCŁAWSKA,
 WYBRZEŻE WYSPIAŃSKIEGO 27, 50-370 WROCŁAW, POLAND

E-mail address: Romuald.Lenczewski@pwr.wroc.pl